

A TOUGHNESS CONDITION FOR FRACTIONAL (k, m) -DELETED GRAPHS REVISITED

WEI GAO^{1,2}, JUAN L.G. GUIRAO³, YAOJUN CHEN¹

ABSTRACT. In computer networks, toughness is an important parameter which is used to measure the vulnerability of the network. Zhou et al. [24] obtains a toughness condition for a graph to be fractional (k, m) -deleted and presents an example to show the sharpness of the toughness bound. In this paper we remark that the previous example does not work and inspired by this fact, we present a new toughness condition for fractional (k, m) -deleted graphs improving the existing one. Finally, we state an open problem.

1. INTRODUCTION

The problem of fractional factor can be discussed as a relaxation of the famous cardinality matching problem which is a crucial problem in operation research. It has wide applications in different areas such as combinatorial polyhedron, scheduling and network design. For instance, certain large data packets are sent to different destinations via several channels in a data transmission network. To finish this work efficiently, it is necessary to divide the large data packets into small ones, and the available assignments of data packets are equal to the problem of fractional flow which can be converted to a problem of fractional factor in a network graph. Some development methods on graph based networks design can refer to Ashwin and Postlethwaite [1], Crouzeilles et al. [5], de Araujo et al. [6], Fardad et al. [7], Haenggi et al. [15], Lanzeni [17], Pishvaei and Rabbani [19], Possani et al. [20], Rahimi and Haghghi [21] and Rizzelli et al. [22] for instance.

All graphs considered in our article are simple (finite, loopless, and without multiple edges). Let $G = (V(G), E(G))$ be a graph with vertex set $V(G)$ and edge set $E(G)$. For any $x \in V(G)$, we denote $d_G(x)$ and $N_G(x)$ by the degree and the neighborhood of x in G , respectively. For

Key words and phrases. Graph, fractional factor, fractional (k, m) -deleted graph, toughness.

2010 Mathematics Subject Classification: 05C70.

$S \subseteq V(G)$, we denote by $G[S]$ the subgraph of G induced by S , and $G - S = G[V(G) \setminus S]$. For two vertex-disjoint subsets S and T of G , we set $e_G(S, T) = |\{e = xy | x \in S, y \in T\}|$ where $|\cdot|$ means cardinality. We denote the minimum degree of G by $\delta(G)$. For more information on the notations and terminologies used in this article, see [3].

Let $k \geq 1$ be an integer and $h : E(G) \rightarrow [0, 1]$ be a function. If $\sum_{x \sim e} h(e) = k$ for any $x \in V(G)$, then we call $G[F_h]$ a *fractional k -factor* of G with indicator function h where $F_h = \{e \in E(G) : h(e) > 0\}$. Zhou [23] introduces the concept of fractional (k, m) -deleted graph, that is, a graph G is called a *fractional (k, m) -deleted graph* if removing any m edges from G , the resulting graph still has a fractional k -factor. A fractional (k, m) -deleted graph is simply called a fractional k -deleted graph if $m = 1$.

The notion of *toughness* was originally introduced by Chvátal [4] in the sense: If G is a complete graph, $t(G) := \infty$. If G is not complete,

$$t(G) := \min\left\{\frac{|S|}{\omega(G - S)} \mid \omega(G - S) \geq 2\right\}$$

and where $\omega(G - S)$ is the number of connected components of $G - S$.

In what follows, we always assume that n is the order of G , i.e., $n = |V(G)|$. Many works have contributed to the analysis of the existence and characterization of fractional factors and fractional (k, m) -deleted graphs. For instance, Liu and Zhang [18] proves that a graph has a fractional k -factor if $t(G) \geq k - \frac{1}{k}$ where $k \geq 2$. Let $\sigma_2(G) := \min\{d_G(u) + d_G(v)\}$ for each pair of non-adjacent vertices u and v in G . Gao and Wang [12] and [13] determine the following sufficient conditions for a graph to be fractional (k, m) -deleted where $k \geq 2$ and $m \geq 0$:

- If $n \geq 4k + 4m - 5$ and $\delta(G) \geq \frac{n}{2}$, then G is a fractional (k, m) -deleted graph.
- If $n \geq 4k + 4m - 3$, $\delta(G) \geq k + m$ and $\max\{d_G(u), d_G(v)\} \geq \frac{n}{2}$ for each pair of non-adjacent vertices u and v of G , then G is a fractional (k, m) -deleted graph.
- If $n \geq \begin{cases} 4k + 4m - 5, & \text{if } (k, m) \neq (3, 0) \\ 8, & \text{if } (k, m) = (3, 0) \end{cases}$, $\delta(G) \geq k + m$ and $\sigma_2(G) \geq n$, then G is a fractional (k, m) -deleted graph.
- If $n \geq 8k + 4m - 7$, $\delta(G) \geq k + m$ and $|N_G(x) \cup N_G(y)| \geq \frac{n}{2}$ for each pair of non-adjacent vertices x, y of G , then G is a fractional (k, m) -deleted graph.

More results on the topic with fractional factor, fractional deleted graphs and other applications can refer to Zhou et al. [25] and [26], Jin [16], Basavanagoud et al. [2], Guirao and Luo [14], Gao et al. [8], and Gao and Wang [9], [10] and [11].

In Zhou et al. [24], the authors proved that a graph G is a fractional (k, m) -deleted graph if $\delta(G) \geq k + 2m$ and $t(G) \geq k + \frac{2m-1}{k}$ for $k \geq 2$. Moreover, they presented an example to show that this toughness bound is “sharp”. We remark that this example does not work to explain the sharpness of the toughness bound determined by Zhou et al. [24] as we show in section 2.

Inspired by this fact we consider that it is still meaningful to discuss the toughness condition for a graph to be fractional (k, m) -deleted, and it is natural to ask whether it would be possible to improve the toughness condition characterization of fractional (k, m) -deleted graph. Thus, in the present paper we shall study the relations between $t(G)$ and the fractional (k, m) -deleted graph. The statement of our main result which improves the Zhou’s et al. previous bound is the following.

Theorem 1. *Let $k \geq 2$ be an integer. A graph G with $\delta(G) \geq k + m$ is a fractional (k, m) -deleted graph if $t(G) \geq k + \frac{m-1}{k}$.*

The structure of this paper is the following. In section 2 we analyze the Zhou’s et al. example and we shall state that it does not work. In section 3 we present the auxiliary results that we shall need for the proof of Theorem 1 which shall be proved in section 4. Finally, in section 5 we shall state an open problem to be study in the future.

2. ON THE ZHOU’S ET AL. EXAMPLE

Now, we shall analyze the sharpness example which is stated in Zhou et al. [24]. Let $k \geq 2$, $n \geq 1$ and $m \geq 0$ be integers. Let K_i be the complete graph of order i and $(j)K_i$ denotes a subgraph of K_i containing j isolated vertices and no edge. The graph of the example is constructed from $K_{n(k-1)}$, $(nk+1)K_{k-1}$ and $K_{(nk+1)(k-1)+2mn}$ as follows:

Let $V((nk+1)K_{k-1}) = \{x_1, x_2, \dots, x_{(nk+1)(k-1)}\}$ and consider the subset $\{y_1, y_2, \dots, y_{(nk+1)(k-1)}\} \subseteq V(K_{(nk+1)(k-1)+2mn})$. $V(G) = V(K_{n(k-1)}) \cup V((nk+1)K_{k-1}) \cup V(K_{(nk+1)(k-1)+2mn})$ and $E(G) = E(K_{n(k-1)}) \cup E((nk+1)K_{k-1}) \cup E(K_{(nk+1)(k-1)+2mn}) \cup \{x_i y_i : i = 1, 2, \dots, (nk+1)(k-1)\} \cup \{u x_i : u \in V(K_{n(k-1)}), i = 1, 2, \dots, (nk+1)(k-1)\}$. It takes $U = (K_{(nk+1)(k-1)+2mn} - \{y_1\}) \cup \{x_1\} \cup K_{n(k-1)}$, and thus $|U| = (nk + n + 1)(k - 1) + 2mn$ and $\omega(G - U) = nk + 2$.

Thus, the authors deduced that $t(G) = \frac{(nk+n+1)(k-1)+2mn}{nk+2}$ which tends to $k + \frac{2m-1}{k}$ when $n \rightarrow \infty$. However, if we consider for the same construction $U := \{y_1, y_2, \dots, y_{(nk+1)(k-1)}\} \cup K_{n(k-1)}$, then we infer $|U| = (nk+n+1)(k-1)$ and $\omega(G-U) = nk+2$. Also, we can check that $\frac{(nk+n+1)(k-1)}{nk+2}$ tends to $k - \frac{1}{k}$ when $n \rightarrow \infty$. It implies that $t(G) < k - \frac{1}{k}$ by the definition of toughness. Therefore, we confirm that this example is not valid to explain the sharpness of the toughness bound determined by Zhou et al. [24].

3. AUXILIARY RESULTS

In this section we present the auxiliary results that we shall need in the proof of our main result.

Lemma 2. (Zhou [23]) *Let $k \geq 1$ and $m \geq 0$ be two integers, let G be a graph and H be a subgraph of G with m edges. Then G is a fractional (k, m) -deleted graph if and only if for any subset S of $V(G)$,*

$$k|S| + \sum_{x \in T} d_{G-S}(x) - k|T| \geq \sum_{x \in T} d_H(x) - e_H(S, T),$$

where $T = \{x : x \in V(G) \setminus S, d_{G-S}(x) - d_H(x) + e_H(x, S) \leq k-1\}$.

We note that Lemma 2 can be easily reformulated in the following equivalent form:

Lemma 3. *Let $k \geq 1$ and $m \geq 0$ be two integers, let G be a graph and H be a subgraph of G with m edges. Then G is a fractional (k, m) -deleted graph if and only if*

$$(1) \quad k|S| + \sum_{x \in T} d_{G-S}(x) - k|T| \geq \sum_{x \in T} d_H(x) - e_H(S, T)$$

for all disjoint subsets S and T of $V(G)$.

Since $\sum_{x \in T} d_H(x) - e_H(S, T) \leq 2m$, we derive the following sufficient condition for fractional (k, m) -deleted graphs.

Lemma 4. *Let $k \geq 1$ and $m \geq 0$ be two integers, let G be a graph and H be a subgraph of G with m edges. Then G is a fractional (k, m) -deleted graph if*

$$k|S| + \sum_{x \in T} d_{G-S}(x) - k|T| \geq 2m$$

for all disjoint subsets S and T of $V(G)$.

The relationship between minimum degree and toughness of graph can be stated as follows.

Lemma 5. (Chvátal [4]) *If a graph G is not complete, then $t(G) \leq \frac{\delta(G)}{2}$.*

The following two lemmas are presented in Liu and Zhang [18] and show the characters of *independent set* (i.e. subsets of the vertex set of G in which any two of them are not adjacent because they reflect the sparsity and stability of the graph system in somehow) and *covering set* (i.e. subsets of the vertex set of G in which each edge has at least one vertex in this set).

Lemma 6. (Liu and Zhang [18]) *Let G be a graph and let $H = G[T]$ such that $\delta(H) \geq 1$ and $1 \leq d_G(x) \leq k - 1$ for every $x \in V(H)$ where $T \subseteq V(G)$ and $k \geq 2$. Let T_1, \dots, T_{k-1} be a partition of the vertices of H satisfying $d_G(x) = j$ for each $x \in T_j$ where we allow some T_j to be empty. If each component of H has a vertex of degree at most $k - 2$ in G , then H has a maximal independent set I and a covering set $C = V(H) - I$ such that*

$$\sum_{j=1}^{k-1} (k-j)c_j \leq \sum_{j=1}^{k-1} (k-2)(k-j)i_j,$$

where $c_j = |C \cap T_j|$ and $i_j = |I \cap T_j|$ for $j = 1, \dots, k - 1$.

Clearly, Lemma 6 also works for $\delta(H) \geq 0$. By the proving process of Lemma 2.2 in [18], we obtain the following Lemma that will be play an important role in the proof of our main result.

Lemma 7. (Liu and Zhang [18]) *Let G be a graph and let $H = G[T]$ such that $d_G(x) = k - 1$ for every $x \in V(H)$ and no component of H is isomorphic to K_k where $T \subseteq V(G)$ and $k \geq 2$. Then there exists an independent set I and a covering set $C = V(H) - I$ of H satisfying*

$$|V(H)| \leq \sum_{i=1}^k (k-i+1)|I^{(i)}| - \frac{|I^{(1)}|}{2}$$

and

$$|C| \leq \sum_{i=1}^k (k-i)|I^{(i)}| - \frac{|I^{(1)}|}{2}$$

where $I^{(i)} = \{x \in I, d_H(x) = k - i\}$ for $1 \leq i \leq k$ and $\sum_{i=1}^k |I^{(i)}| = |I|$.

Note that for each vertex $x \in I$, $d_H(x) = k - 1$ and there exists a vertex $y \in I - \{x\}$ such that $N_H(x) \cap N_H(y) \neq \emptyset$.

4. PROOF OF THEOREM 1

If G is complete, clearly G is a fractional (k, m) -deleted graph by means of $\delta(G) \geq k + m$. In the sequel, we assume that G is not complete. Suppose that G satisfies the hypothesis of Theorem 1, but is not a fractional (k, m) -deleted graph. By Lemma 4 and the fact that $\sum_{x \in T} d_H(x) - e_H(T, S) \leq 2m$ for all S, T disjoint subsets of $V(G)$ and $H \subseteq E(G)$, there exist disjoint subsets S and T of $V(G)$ such that

$$(2) \quad k|S| + d_{G-S}(T) - k|T| \leq 2m - 1.$$

We select S and T such that $|T|$ is minimum. If $T = \emptyset$, then $\sum_{x \in T} d_H(x) - e_H(T, S) = 0$ in view of its definition. Using Lemma 3, (1) becomes $k|S| \geq 0$ and thus G is a fractional (k, m) -deleted graph, which is a contradiction. Hence, we have $T \neq \emptyset$.

If there exists $x \in T$ satisfying $d_{G-S}(x) \geq k$, then the subsets S and $T \setminus \{x\}$ satisfy (2) as well. This contradicts the selection rule of S and T . It implies that $d_{G-S}(x) \leq k - 1$ for any $x \in T$. Furthermore, we infer that $S \neq \emptyset$.

Let l be the number of the components of $H' = G[T]$ which are isomorphic to K_k and let $T_0 = \{x \in V(H') | d_{G-S} = 0\}$. Let H be the subgraph obtained from $H' - T_0$ by deleting those l components isomorphic to K_k .

If $|V(H)| = 0$, then from (2) we obtain $1 \leq |S| \leq |T_0| + l + \frac{2m-1}{k}$. Clearly, $|T_0| + l \geq 1$ since $|T| \neq 0$. If $|T_0| + l = 1$, then $|S| \leq 1 + \frac{2m-1}{k}$ and $2k + \frac{2m-2}{k} \leq 2t(G) \leq \delta(G) \leq k - 1 + |S| \leq k + \frac{2m-1}{k}$ which contradicts to $k \geq 2$. Hence $\omega(G - S) \geq |T_0| + l \geq 2$ and $t(G) \leq \frac{|S|}{\omega(G-S)} \leq \frac{|S|}{|T_0|+l} \leq \frac{1}{2} + \frac{2m-1}{2k}$. This contradicts $t(G) \geq k + \frac{m-1}{k}$ and $k \geq 2$. Therefore, we have $|V(H)| > 0$.

Let $H = H_1 \cup H_2$ where H_1 is the union of components of H which satisfies that $d_{G-S}(x) = k - 1$ for every vertex $x \in V(H_1)$ and $H_2 = H - H_1$. By Lemma 7, H_1 has a maximum independent set I_1 and the covering set $C_1 = V(H_1) - I_1$ such that

$$(3) \quad |V(H_1)| \leq \sum_{i=1}^k (k - i + 1) |I^{(i)}| - \frac{|I^{(1)}|}{2},$$

and

$$(4) \quad |C_1| \leq \sum_{i=1}^k (k - i) |I^{(i)}| - \frac{|I^{(1)}|}{2},$$

where $I^{(i)} = \{x \in I_1, d_{H_1}(x) = k - i\}$ for $1 \leq i \leq k$ and $\sum_{i=1}^k |I^{(i)}| = |I_1|$.

On the other hand, let $T_j = \{x \in V(H_2) | d_{G-S}(x) = j\}$ for $1 \leq j \leq k - 1$. By the definition of H and H_2 we can also see that each component of H_2 has a vertex of degree at most $k - 2$ in $G - S$. According to Lemma 6, H_2 has a maximal independent set I_2 and the covering set $C_2 = V(H_2) - I_2$ such that

$$(5) \quad \sum_{j=1}^{k-1} (k-j)c_j \leq \sum_{j=1}^{k-1} (k-2)(k-j)i_j,$$

where $c_j = |C_2 \cap T_j|$ and $i_j = |I_2 \cap T_j|$ for every $j = 1, \dots, k - 1$. Set $W = V(G) - S - T$ and $U = S \cup C_1 \cup (N_G(I_1) \cap W) \cup C_2 \cup (N_G(I_2) \cap W)$. In light of computation, we infer

$$\begin{aligned} & |C_2| + |N_G(I_2) \cap W| = |V(H_2)| - |I_2| + |N_{G-S-T}(I_2)| \\ & = |V(H_2)| - |I_2| + |N_{G-S}(I_2)| - |N_T(I_2)| \\ & = (|V(H_2)| - |I_2| - |N_T(I_2)|) + |N_{G-S}(I_2)| \\ & \leq (|V(H_2)| - |I_2| - |N_{H_2}(I_2)|) + |N_{G-S}(I_2)| \\ & \leq 0 + \sum_{j=1}^{k-1} j i_j = \sum_{j=1}^{k-1} j i_j. \end{aligned}$$

Furthermore, we get

$$(6) \quad |U| \leq |S| + |C_1| + \sum_{j=1}^{k-1} j i_j + \sum_{i=1}^k (i-1) |I^{(i)}|$$

and

$$(7) \quad \omega(G - S) \geq t_0 + l + |I_1| + \sum_{j=1}^{k-1} i_j,$$

where $t_0 = |T_0|$. Let $t(G) = t$. Then when $\omega(G - S) > 1$, we have

$$(8) \quad |U| \geq t\omega(G - S),$$

and it is also held when $\omega(G - S) = 1$ (in this case, $t\omega(G - S) = t \leq \frac{\delta(G)}{2} \leq \frac{|S| + d_{G-S}(x)}{2} \leq \frac{|U|}{2}$ where $x \in T$).

By (6)-(8), we yield

$$(9) \quad |S| + |C_1| \geq \sum_{j=1}^{k-1} (t-j)i_j + t(t_0 + l) + t|I_1| - \sum_{i=1}^k (i-1)|I^{(i)}|.$$

Note that $|T| = t_0 + lk + |V(H_1)| + |V(H_2)|$, and

$$\begin{aligned}
& k|T| - d_{G-S}(T) = kt_0 + lk^2 + k|V(H_1)| + k|V(H_2)| - d_{G-S}(T) \\
& = kt_0 + lk^2 + k|V(H_1)| + k|V(H_2)| - lk(k-1) - (k-1)|V(H_1)| \\
& \quad - \sum_{j=1}^{k-1} j(i_j + c_j) \\
& = kt_0 + lk + |V(H_1)| + \sum_{j=1}^{k-1} (k-j)i_j + \sum_{j=1}^{k-1} (k-j)c_j.
\end{aligned}$$

Thus, from (2) we have

$$kt_0 + kl + |V(H_1)| + \sum_{j=1}^{k-1} (k-j)i_j + \sum_{j=1}^{k-1} (k-j)c_j \geq k|S| - 2m + 1.$$

Combining with (9) we have

$$\begin{aligned}
& kt_0 + kl + |V(H_1)| + \sum_{j=1}^{k-1} (k-j)i_j + \sum_{j=1}^{k-1} (k-j)c_j + k|C_1| \\
& \geq \sum_{j=1}^{k-1} (kt - kj)i_j + kt(t_0 + l) + kt|I_1| - k \sum_{i=1}^k (i-1)|I^{(i)}| - 2m + 1.
\end{aligned}$$

Thus

$$\begin{aligned}
(10) \quad & |V(H_1)| + \sum_{j=1}^{k-1} (k-j)c_j + k|C_1| \\
& \geq \sum_{j=1}^{k-1} (kt - kj - k + j)i_j + k(t-1)(t_0 + l) + kt|I_1| \\
(11) \quad & -k \sum_{i=1}^k (i-1)|I^{(i)}| - 2m + 1.
\end{aligned}$$

By (3) and (4),

$$(12) \quad |V(H_1)| + k|C_1| \leq \sum_{i=1}^k (k^2 - (i-1)k - (i-1))|I^{(i)}| - \frac{(k+1)|I^{(1)}|}{2}.$$

Using (5),(10) and (12), we have

$$\begin{aligned}
(13) \quad & \sum_{j=1}^{k-1} (k-2)(k-j)i_j + \sum_{i=1}^k (k^2 - (i-1)k - (i-1))|I^{(i)}| \\
& \geq \sum_{j=1}^{k-1} (kt - kj - k + j)i_j + k(t-1)(t_0 + l) + kt|I_1| + \frac{(k+1)|I^{(1)}|}{2} \\
& \quad - k \sum_{i=1}^k (i-1)|I^{(i)}| - 2m + 1.
\end{aligned}$$

Now, we shall discuss the following cases according to the value of $t_0 + l$.

Case 1. $t_0 + l \geq 1$. In this case, by (13) and $k(t-1)(t_0 + l) - 2m + 1 \geq k(k + \frac{m-1}{k} - 1) - 2m + 1 = k^2 - m - 1$,

$$\begin{aligned}
(14) \quad & \sum_{j=1}^{k-1} (k-2)(k-j)i_j + \sum_{i=1}^k (k^2 - (i-1)k - (i-1))|I^{(i)}| \\
& \geq \sum_{j=1}^{k-1} (kt - kj - k + j)i_j + kt|I_1| + \frac{(k+1)|I^{(1)}|}{2} - k \sum_{i=1}^k (i-1)|I^{(i)}| \\
& \quad + k^2 - m - 1.
\end{aligned}$$

Claim 8. If $t_0 + l \geq 1$, then $|I_2| \neq 0$.

Proof. Suppose $|I_2| = 0$. Then $|I_1| \neq 0$ by $|V(H)| > 0$ and (14) becomes

$$(15) \quad (|I_1| - 1)k^2 - (t|I_1| + \frac{|I^{(1)}|}{2})k - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} + m + 1 \geq 0.$$

In light of (15) and $t \geq k + \frac{m-1}{k}$ we have

$$\begin{aligned}
0 & \leq (|I_1| - 1)k^2 - (t|I_1| + \frac{|I^{(1)}|}{2})k - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} + m + 1 \\
& \leq (|I_1| - 1)k^2 - ((k + \frac{m-1}{k})|I_1| + \frac{|I^{(1)}|}{2})k - \sum_{i=1}^k (i-1)|I^{(i)}| \\
& \quad - \frac{|I^{(1)}|}{2} + m + 1 \\
& = -k^2 - \frac{|I^{(1)}|}{2}k - (m-1)|I_1| - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} + m + 1 < 0,
\end{aligned}$$

a contradiction. The last step follows from $|I_1| \geq 1$ and $k \geq 2$. \square

Claim 9. *If $t_0 + l \geq 1$, then $|I_1| \neq 0$.*

Proof. Suppose $|I_1| = 0$. We yield $|I_2| \neq 0$ by $|V(H)| > 0$ and (14) becomes

$$\sum_{j=1}^{k-1} (k-2)(k-j)i_j \geq \sum_{j=1}^{k-1} (kt - kj - k + j)i_j + k^2 - m - 1.$$

Note that

$$\begin{aligned} & (k-2)(k-j) - kt + kj + k - j \\ & \leq (k-2)(k-j) - k\left(k + \frac{m-1}{k}\right) + kj + k - j \\ & = -k + j - m + 1 \leq -m. \end{aligned}$$

We have

$$\sum_{j=1}^{k-1} (k-2)(k-j)i_j - \sum_{j=1}^{k-1} (kt - kj - k + j)i_j - k^2 + m + 1 < 0$$

in view of $k \geq 2$. \square

From Claim 8 and Claim 9, we can see that $|I_1| > 0$ and $|I_2| > 0$. According to $|I_2| \geq 1$ and

$$\sum_{j=1}^{k-1} (k-2)(k-j)i_j - \sum_{j=1}^{k-1} (kt - kj - k + j)i_j + m \leq 0,$$

we have

$$\begin{aligned} & \sum_{i=1}^k (k^2 - (i-1)k - (i-1))|I^{(i)}| \\ & \geq kt|I_1| + \frac{(k+1)|I^{(1)}|}{2} - k \sum_{i=1}^k (i-1)|I^{(i)}| + k^2 - 1. \end{aligned}$$

That is to say,

$$(16) \quad (|I_1| - 1)k^2 - (t|I_1| + \frac{|I^{(1)}|}{2})k - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} + 1 \geq 0.$$

In light of (16) and $t \geq k + \frac{m-1}{k}$ we have

$$\begin{aligned}
0 &\leq (|I_1| - 1)k^2 - (t|I_1| + \frac{|I^{(1)}|}{2})k - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} + 1 \\
&\leq (|I_1| - 1)k^2 - ((k + \frac{m-1}{k})|I_1| + \frac{|I^{(1)}|}{2})k - \sum_{i=1}^k (i-1)|I^{(i)}| \\
&\quad - \frac{|I^{(1)}|}{2} + 1 \\
&= -k^2 - \frac{|I^{(1)}|}{2}k - (m-1)|I_1| - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} + 1 < 0,
\end{aligned}$$

a contradiction.

Case 2. $t_0 + l = 0$. In this case, by (13) we have,

$$\begin{aligned}
(17) \quad &\sum_{j=1}^{k-1} (k-2)(k-j)i_j + \sum_{i=1}^k (k^2 - (i-1)k - (i-1))|I^{(i)}| \\
&\geq \sum_{j=1}^{k-1} (kt - kj - k + j)i_j + kt|I_1| + \frac{(k+1)|I^{(1)}|}{2} - k \sum_{i=1}^k (i-1)|I^{(i)}| \\
&\quad - 2m + 1.
\end{aligned}$$

Claim 10. *If $t_0 + l = 0$, then $|I_2| \neq 0$.*

Proof. Suppose $|I_2| = 0$. Then $|I_1| \neq 0$ by $|V(H)| > 0$, $|V(T)| = |V(H_1)|$ and $k|S| \leq k|T| - d_{G-S}(T) + 2m - 1 = |T| + 2m - 1$. If $|I_1| \leq k$, then $|T| \leq k^2$ and $|S| \leq \frac{|T| + 2m - 1}{k} \leq k + \frac{2m-1}{k}$. Then $2k + \frac{2m-2}{k} \leq 2t \leq \delta(G) \leq |S| + (k-1) \leq 2k - 1 + \frac{2m-1}{k}$, which contradicts to $k \geq 2$. Hence, we have $|I_1| \geq k + 1$.

Using (17), we have

$$(18) \quad |I_1|k^2 - (t|I_1| + \frac{|I^{(1)}|}{2})k - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} + 2m - 1 \geq 0.$$

In light of (18), $t \geq k + \frac{m-1}{k} |I_1| \geq k + 1$ and $k \geq 2$, we get

$$\begin{aligned}
0 &\leq |I_1|k^2 - (t|I_1| + \frac{|I^{(1)}|}{2})k - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} + 2m - 1 \\
&\leq |I_1|k^2 - ((k + \frac{m-1}{k})|I_1| + \frac{|I^{(1)}|}{2})k - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} \\
&\quad + 2m - 1 \\
&= -\frac{|I^{(1)}|}{2}k - (m-1)|I_1| - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} + 2m - 1 < 0,
\end{aligned}$$

a contradiction. \square

Claim 11. *If $t_0 + l = 0$, then $|I_1| \neq 0$.*

Proof. Suppose $|I_1| = 0$. Then $|I_2| \neq 0$ using $|V(H)| > 0$. In terms of (17), we infer

$$\sum_{j=1}^{k-1} (k-2)(k-j)i_j \geq \sum_{j=1}^{k-1} (kt - kj - k + j)i_j - 2m + 1.$$

Note that

$$\max\{(k-2)(k-j) - kt + kj + k - j\} = -m.$$

Since $|C_2| \leq (k-2) + (|I_1| - 1)(k-1-1) = |I_2|(k-2)$ and $|T| \leq |I_2|(k-1)$, we get $|S| \leq \frac{|T| + 2m - 1}{k} \leq |I_2| + \frac{2m-1-|I_2|}{k}$. If $|I_2| = 1$, then $|S| \leq 1 + \frac{2m-2}{k}$ and $2k + \frac{2m-2}{k} \leq 2t \leq \delta(G) \leq |S| + (k-1) = k + \frac{2m-2}{k}$, a contradiction. Hence, we get $|I_2| \geq 2$ and

$$\sum_{j=1}^{k-1} (k-2)(k-j)i_j - \sum_{j=1}^{k-1} (kt - kj - k + j)i_j + 2m - 1 < 0.$$

This implies that $|I_1| \neq 0$. \square

From Claim 10 and Claim 11, we can see that $|I_1| > 0$ and $|I_2| > 0$. In terms of $|I_2| \geq 1$, we deduce $\sum_{j=1}^{k-1} (kt - kj - k + j)i_j \geq \sum_{j=1}^{k-1} (k-2)(k-j)i_j + m$.

Thus, we obtain

$$\begin{aligned}
&\sum_{i=1}^k (k^2 - (i-1)k - (i-1))|I^{(i)}| \\
&\geq kt|I_1| + \frac{(k+1)|I^{(1)}|}{2} - k \sum_{i=1}^k (i-1)|I^{(i)}| - m + 1,
\end{aligned}$$

which reveals

$$(19) \quad |I_1|k^2 - (t|I_1| + \frac{|I^{(1)}|}{2})k - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} + m - 1 \geq 0.$$

In light of (19) $|I_1| \geq 1$ and $t \geq k + \frac{m-1}{k}$, we have

$$\begin{aligned} 0 &\leq |I_1|k^2 - (t|I_1| + \frac{|I^{(1)}|}{2})k - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} + m - 1 \\ &\leq |I_1|k^2 - ((k + \frac{m-1}{k})|I_1| + \frac{|I^{(1)}|}{2})k - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} \\ &\quad + m - 1 \\ &= -\frac{|I^{(1)}|}{2}k - (m-1)|I_1| - \sum_{i=1}^k (i-1)|I^{(i)}| - \frac{|I^{(1)}|}{2} + m - 1 < 0. \end{aligned}$$

Therefore the proof of Theorem 1 is over. ■

5. CONCLUSIONS

From the detailed proof of Theorem 1 presented in the previous section, we see that the condition $\delta(G) \geq k + m$ is only used to ensure that G is a fractional (k, m) -deleted graph when G is complete. If we assume that G is not complete in our main result, then the condition $\delta(G) \geq k + m$ can be deleted.

We believe that the toughness bound presented in our paper is still not sharp, although it improves the previous existing result. Therefore, we give the following conjecture on the relationship between fractional (k, m) -deleted graph and toughness.

Conjecture 12. *If $t(G) \geq k - \frac{1}{k}$ and $k \geq \Theta(m)$ (here $\Theta(m)$ is a function of m and $\Theta(m) \sim O(m)$), then G is a fractional (k, m) -deleted graph.*

6. ACKNOWLEDGMENTS

This work has been partially supported by MINECO grant number MTM2014-51891-P and Fundación Séneca de la Región de Murcia grant number 19219/PI/14 and National Science Foundation of China grant number 11401519.

REFERENCES

- [1] P. ASHWIN, C. POSTLETHWAITE, *On designing heteroclinic networks from graphs*, *Physica D* **265** (2013), 26–39.
- [2] B. BASAVANAGOUD, V. R. DESAI, S. PATIL, (β, α) -Connectivity index of graphs, *Appl. Math. Nonl. Sc.* **2** (2017), 21–30.
- [3] J. A. BONDY, U. S. R. MUTRY, *Graph Theory*, Springer, Berlin, 2008.
- [4] V. CHVÁTAL, *Tough graphs and hamiltonian circuits*, *Discrete Math.* **5** (1973), 215–228.
- [5] R. CROUZEILLES, M.L. LORINI, C.E.D. GRELE, *Applying graph theory to design networks of protected areas: using inter-patch distance for regional conservation planning*, *Nat. Conservacao*, **9** (2011), 219–224.
- [6] D.R.B. DE ARAUJO, J. F. MARTINS, C. J. A. BASTOS, *New graph model to design optical networks*, *IEEE Commun. Lett.* **19** (2015), 2130–2133.
- [7] M. FARDAD, F. LIN, M. R. JOVANOVIC, *Design of optimal sparse interconnection graphs for synchronization of oscillator networks*, *IEEE T. Automat. Contr.* **59** (2014), 2457–2462.
- [8] W. GAO, Y. GUO, K. Y. WANG, *Ontology algorithm using singular value decomposition and applied in multidisciplinary*, *Cluster Comput.* **19** (2016), 2201–2210.
- [9] W. GAO, W. F. WANG, *New isolated toughness condition for fractional (g, f, n) -critical graphs*, *Colloq. Math.* **147** (2017), 55–66.
- [10] W. GAO, W. F. WANG, *The eccentric connectivity polynomial of two classes of nanotubes*, *Chaos Soliton & Fractals* **89** (2016), 290–294.
- [11] W. GAO, W. F. WANG, *The fifth geometric arithmetic index of bridge graph and carbon nanocones*, *J. Differ. Equ. Appl.* 2016, <http://dx.doi.org/10.1080/10236198.2016.1197214>.
- [12] W. GAO, W. F. WANG, *Degree conditions for fractional (k, m) -deleted graphs*, *Ars Combin.* **113A** (2014), 273–285.
- [13] W. GAO, W. F. WANG, *A neighborhood union condition for fractional (k, m) -deleted graphs*, *Ars Combin.* **113A** (2014), 225–233.
- [14] J.L.G. GUIRAO, A.C.J. LUO, *New trends in nonlinear dynamics and chaoticity*, *Nonlinear Dynam.* **84** (2016), 1–2.
- [15] M. HAENGGI, J. ANDREWS, F. BACCELLI, O. DOUSSE, M. FRANCESCHETTI, D. TOWSLEY, *Stochastic geometry and random graphs for the analysis and design of wireless networks*, *IEEE J. Sel. Area. Comm.* **27** (2009), 1025–1028.
- [16] J. H. JIN, *Multiple solutions of the Kirchhoff-type problem in R^N* , *Appl. Math. Nonl. Sc.* **1** (2016), 229–238.
- [17] S. LANZENI, E. MESSINA, F. ARCHETTI, *Graph models and mathematical programming in biochemical network analysis and metabolic engineering design*, *Comput. Math. Appl.* **55** (2008), 970–983.
- [18] G.Z. LIU, L. J. ZHANG, *Toughness and the existence of fractional k -factors of graphs*, *Discrete Math.* **308** (2008), 1741–1748.
- [19] M. S. PISHVAEE, M. RABBANI, *A graph theoretic-based heuristic algorithm for responsive supply chain network design with direct and indirect shipment*, *Adv. Eng. Softw.* **42** (2011), 57–63.
- [20] V.N. POSSANI, V. CALLEGARO, A.I. REIS, R.P. RIBAS, F.D. MARQUES, L.S. DA ROSA, *Graph-based transistor network generation method for supergate design*, *IEEE T. VLSI Syst.* **24** (2016), 692–705.

- [21] M. RAHIMI, A. HAGHIGHI, *A graph portioning approach for hydraulic analysis-design of looped pipe networks*, Water Resour. Manag. **29** (2015), 5339–5352.
- [22] G. RIZZELLI, M. TORNATORE, G. MAIER, A. PATTAVINA, *Impairment-aware design of translucent DWDM networks based on the k -path connectivity graph*, J. Opt. Commun. Netw. **4** (2012), 356–365.
- [23] S. ZHOU, *A neighborhood condition for graphs to be fractional (k, m) -deleted graphs*, Glasg. Math. J., **52(1)** (2010), 33–40.
- [24] S.Z. ZHOU, Z.R. SUN, H. YE, *A toughness condition for fractional (k, m) -deleted graphs*, Inform. Process. Lett. **113** (2013), 255–259.
- [25] S.Z. ZHOU, L. XU, Y. XU, *A sufficient condition for the existence of a k -factor excluding a given r -factor*, Appl. Math. Nonl. Sc. **2** (2017), 13–20.
- [26] S. ZHOU, F. YANG, Z. R. SUN, *A neighborhood condition for fractional ID - $[a, b]$ -factor-critical graphs*, Discussiones Mathematicae Graph Theory **36(2)** (2016), 409–418.

¹ DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, CHINA

E-mail address: gaowei@ynnu.edu.cn

² SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, YUNNAN NORMAL UNIVERSITY, KUNMING 650500, CHINA

³ DEPARTAMENTO DE MATEMÁTICA APLICADA Y ESTADÍSTICA. UNIVERSIDAD POLITÉCNICA DE CARTAGENA, HOSPITAL DE MARINA, 30203-CARTAGENA, REGIÓN DE MURCIA, SPAIN—CORRESPONDING AUTHOR—

E-mail address: juan.garcia@upct.es