

# NEW AND MODIFIED ECCENTRIC INDICES OF OCTAGONAL GRID $O_n^m$

M. NAEEM<sup>1</sup>, M. K. SIDDIQUI<sup>1</sup>, J. L. G. GUIRAO<sup>2</sup> AND W. GAO<sup>3</sup>

ABSTRACT. The eccentricity  $\varepsilon_u$  of vertex  $u$  in a connected graph  $G$ , is the distance between  $u$  and a vertex farthest from  $u$ . The aim of the present paper is to introduce new eccentricity based index and eccentricity based polynomial, namely modified augmented eccentric connectivity index and modified augmented eccentric connectivity polynomial respectively. As an application we compute these new indices for octagonal grid  $O_n^m$  and we compare the results obtained with the ones obtained by other indices like Ediz eccentric connectivity index, modified eccentric connectivity index and modified eccentric connectivity polynomial  $ECP(G, x)$ .

## 1. INTRODUCTION

In recent years graph theory is extensively used in the branch of mathematical chemistry and some people call it as *chemical graph theory* because this theory is related with the practical applications of graph theory for solving the molecular problems. In mathematics a model of chemical system portrays a chemical graph that deals to explain the relations between its segments such as its atoms, bonds between atoms, cluster of atoms or molecules.

A *connected simple graph*  $G = (V(G) \cup E(G))$  is a graph consisting of  $n$  vertices ( $V(G)$ ) and  $m$  edges ( $E(G)$ ) in which there is path between any of two its vertices. A *network* is merely a connected graph consisting of no multiple edges and loops. The *degree of a vertex*  $v$  in  $G$  is the number of edges which are incident to the vertex  $v$  and will be represented by  $d_v$ . In a graph  $G$ , if there is no repetition of vertices in  $(u - v)$  walk then such kind of walk is called  $(u - v)$  *path*. The number of edges in  $(u - v)$  path is called its *length*. The *distance*  $d(u, v)$  from vertex  $u$  to vertex  $v$  is the length of a shortest  $(u - v)$  path in a graph  $G$  where  $u, v \in G$ . In a connected graph  $G$ , the *eccentricity*  $\varepsilon_v$  of a vertex  $v$  is the distance between  $v$  and a vertex furthest from  $v$  in  $G$ . Thus,  $\varepsilon_v = \max_{v \in V(G)} d(v, u)$ . Therefore the maximum eccentricity over all vertices of  $G$  is the *diameter* of  $G$  which is denoted by  $D(G)$ .

A graph can be recognized by a different type of numeric number, a polynomial, a sequence of numbers or a matrix. A *topological index* is a numeric quantity that is

---

2010 *Mathematics Subject Classification*. Primary: 05C12, 05C90.

*Key words and phrases*. Degree, Eccentricity, Ediz eccentric connectivity index, modified eccentric connectivity index, modified eccentric connectivity polynomial  $ECP(G, x)$ , octagonal grid  $O_n^m$ .

associated with a graph which characterize the topology of graph and is invariant under graph automorphism. Over the years topological indices like Wiener index Balabans index [24, 25, 26], Hosoya index [16, 17], Randić index [19] and so on, have been studied extensively and recently the research and interest in this area has been increased exponentially. See too for more information [3, 13, 14, 18, 21, 23].

There are some major classes of topological indices such as *distance based topological indices*, *eccentricity based topological indices*, *degree based topological indices* and *counting related polynomials* and indices of graphs. In this article we shall consider the eccentricity based indices. We note that in [5] is introduced the *total eccentricity* of a graph  $G$  and is defined as the sum of eccentricities of all vertices of a given graph  $G$  and denote by  $\zeta(G)$ . It is easy to see that for a  $k$ -regular graph  $G$  is held  $\zeta(G) = k\zeta(G)$ .

The *Eccentric-connectivity index*  $\xi(G)$  which was proposed by Sharma, Goswami and Madan defined as [20]:

$$(1) \quad \xi(G) = \sum_{u \in V(G)} d_u \epsilon_u,$$

Another very relevant and special eccentricity based topological index is *connective Eccentric index*  $C^\xi(G)$  that was proposed by Gupta et al. in [11]. The *connective eccentric index* is defined as.

$$(2) \quad C^\xi(G) = \sum_{u \in V(G)} \frac{d_u}{\epsilon_u},$$

In 2010, A. R. Ashrafi and M. Ghorbani [1] introduces the so called *modified eccentric connectivity index*  $\xi_c(G)$  and it is defined as

$$(3) \quad \xi_c(G) = \sum_{v \in V(G)} (S_v \epsilon_v),$$

where  $S_v = \sum_{u \in N(v)} d_u$  that is  $S_v$  is the sum of degrees of all vertices adjacent to vertex  $v$ .

In 2010, S. Ediz et al., [8], defined *Ediz eccentric connectivity index* of  $G$  as

$$(4) \quad {}^E\xi^c(G) = \sum_{v \in V(G)} \left( \frac{S_v}{\epsilon_v} \right),$$

Similar to other topological polynomials, the corresponding polynomial, that is, the *modified eccentric connectivity polynomial* of a graph, is defined as, [6]:

$$(5) \quad \xi_c(G, x) = \sum_{u \in V(G)} S_u x^{\epsilon_u},$$

so that the *modified eccentric connectivity index* is the first derivative of this polynomial for  $x = 1$ .

Motivated by these above eccentricity indices, in this article we introduce what we call *modified augmented eccentric connectivity index*  $^{MA}\xi(G)$ , as

$$(6) \quad ^{MA}\xi^c(G) = \sum_{v \in V(G)} (M_v \epsilon_v),$$

where  $M_v = \prod_{u \in N(v)} d_u$  that is denotes the product of degrees of all neighbors of vertex  $v$  of  $G$ .

In the same way, we define the *modified augmented eccentric connectivity polynomial*  $^{MA}\xi^c(G, x)$ , as

$$(7) \quad ^{MA}\xi^c(G, x) = \sum_{v \in V(G)} M_v x^{\epsilon_v}$$

For more information and properties of eccentricity based topological index, see for instance [2, 7, 9, 10, 12, 15, 27].

The aim of this paper is is the introduction of the augmented eccentric connectivity index and modified augmented eccentric connectivity polynomial. As an application we shall compute these new indices for octagonal grid  $O_n^m$  and we shall compare the results obtained with the ones obtained by other indices like Ediz eccentric connectivity index, modified eccentric connectivity index and modified eccentric connectivity polynomial  $ECP(G, x)$  via their computation too.


## 2. OCTAGONAL GRID $O_n^m$

In [4] and [22] Diudea et al. constructed a  $C_4C_8$  net as a trivalent decoration made by alternating squares  $C_4$  and octagons  $C_8$  in two different ways. One is by alternating squares  $C_4$  and octagons  $C_8$  in different ways denoted by  $C_4C_8(S)$  and other is by alternating rhombus and octagons in different ways denoted by  $C_4C_8(R)$ . We denote  $C_4C_8(R)$  by  $O_n^m$  see Figure 1. In [21] they also called it as *the Octagonal grid*.

For  $n, m \geq 2$  the Octagonal grid  $O_n^m$ , is the grid with  $m$  rows and  $n$  columns of octagons. The symbols  $V(O_n^m)$  and  $E(O_n^m)$  will denote the vertex set and the edge set of  $O_n^m$ , respectively.

$$\begin{aligned} V(O_n^m) &= \{u_s^t : 1 \leq s \leq n, 1 \leq t \leq m+1\} \cup \{v_s^t : 1 \leq s \leq n; 1 \leq t \leq m+1\} \\ &\cup \{w_s^t : 1 \leq s \leq n+1, 1 \leq t \leq m\} \cup \{y_s^t : 1 \leq s \leq n+1, 1 \leq t \leq m\}. \end{aligned}$$

$$\begin{aligned} E(O_n^m) &= \{u_s^t v_s^t : 1 \leq s \leq n, 1 \leq t \leq m+1\} \cup \{u_s^t w_s^t; 1 \leq s \leq n, 1 \leq t \leq m\} \\ &\cup \{w_s^t y_s^t : 1 \leq s \leq n+1, 1 \leq t \leq m\} \cup \{v_s^t w_{s+1}^t : 1 \leq s \leq n, 1 \leq t \leq m\} \\ &\cup \{v_s^t y_{s+1}^{t-1} : 1 \leq s \leq n, 2 \leq t \leq m+1\} \cup \{u_s^{t+1} y_s^t : 1 \leq s \leq n, 2 \leq t \leq m\}. \end{aligned}$$

FIGURE 1. The Octagonal grid  $O_n^m$ .

In this paper, we consider  $O_n^m$  with  $n = m$ .

### 3. STATEMENT OF MAIN RESULTS

As we have said previously for  $O_n^m$  with  $n = m$  we shall compute modified eccentric connective index, Ediz eccentric connectivity index, modified eccentric connective polynomial, modified augmented eccentric connective index and modified augmented eccentric connective polynomial and we shall compare the results obtained. For this we have discussed two cases of  $n$ , when  $n \equiv 0 \pmod{2}$  and when  $n \equiv 1 \pmod{2}$ . Also to avoid any ambiguity related to Figure 1 note that the vertices  $u_s^t = u_s^t$ .

**Theorem 1.** For every  $n \geq 4$  and  $n \equiv 0 \pmod{2}$  consider the graph of  $G \cong O_n^m$ , with  $n = m$ . Then the modified eccentric connectivity index  $\xi_c(G)$  of  $G$  is equal to

$$\begin{aligned}
\xi_c(O_n^m) &= 225n^2 - 112n + 28 \\
&+ 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} \{4n - 3(s-1) - t\} \right] + 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s \{4(n+1) - s - 3t\} \right] \\
&+ 36 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} \{3(n-t) + s + 2\} \right] + 36 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} \{n + 3s - t - 1\} \right] \\
&+ 36 \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} \{3(n-s) + t + 1\} \right] + 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n \{n - s + 3t - 2\} \right] \\
&+ 36 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s \{4s + t - n - 3\} \right] + 36 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n \{s + 3t - 4\} \right].
\end{aligned}$$

*Proof.* Let  $G$  be the graph of  $O_n^m$ . Note that graph of  $O_n^m$  is a symmetric about reflection and rotation at right angles. Thus the eccentricities  $\varepsilon_{u_s^t} = \varepsilon_{v_{n+1-s}^t}$  and from the symmetry at right angles we can obtain that the eccentricities  $\varepsilon_{y_s^t} = \varepsilon_{u_t^s}$ ,  $\varepsilon_{w_s^t} = \varepsilon_{v_t^s}$ . Therefore, from Table 1 and formula (3), given below, the modified eccentric connectivity index  $\xi_c(G)$  of  $O_n^m$  is equal to

$$\xi_c(G) = \sum_{v \in V(G)} (S_v \varepsilon_v) = 4 \sum_{u_s^t \in V(G)} (S_{u_s^t} \varepsilon_{u_s^t}),$$

$$\begin{aligned}
\xi_c(O_n^m) &= 4 \left[ 2 \times 4 \times 4n + 2 \sum_{s=2}^{\frac{n}{2}+1} 5\{4n+1-s\} + 2 \sum_{s=\frac{n}{2}+2}^n 5\{3n+s-1\} \right] \\
&+ 4 \left[ \sum_{t=2}^{\frac{n}{2}+1} 7(4n-t) + \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} 9\{4n-3(s-1)-t\} \right] \right] \\
&+ \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s 9\{4(n+1)-s-3t\} \right] + \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} 9\{3(n-t)+s+2\} \right] \\
&+ \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} 9\{n+3s-t-1\} \right] + \sum_{t=\frac{n}{2}+2}^n 7\{3(n-1)+t+1\} \\
&+ \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} 9\{3(n-s)+t+1\} \right] + \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n 9\{n-s+3t-2\} \right] \\
&+ \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s 9\{4s+t-n-3\} \right] + \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n 9\{s+3t-4\} \right].
\end{aligned}$$

After some easy calculations we get

$$\begin{aligned}
\xi_c(O_n^m) &= 225n^2 - 112n + 28 \\
&+ 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} \{4n-3(s-1)-t\} \right] + 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s \{4(n+1)-s-3t\} \right] \\
&+ 36 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} \{3(n-t)+s+2\} \right] + 36 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} \{n+3s-t-1\} \right] \\
&+ 36 \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} \{3(n-s)+t+1\} \right] + 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n \{n-s+3t-2\} \right] \\
&+ 36 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s \{4s+t-n-3\} \right] + 36 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n \{s+3t-4\} \right].
\end{aligned}$$

□

TABLE 1. Partition of vertices of the type  $u_s^t$  of  $O_n^m$  based on degree sum and eccentricity of each vertex when  $n \equiv 0 \pmod{2}$ .

Representative	$S_{u_s^t}$	eccentricity	Range	Frequency
$u_s^t$	4	$4n - s + 1$	$t = 1, n + 1; s = 1$	2
$u_s^t$	5	$4n - s + 1$	$t = 1, n + 1,$ $2 \leq s \leq \frac{n}{2} + 1$	$n$
$u_s^t$	5	$3n + s - 1$	$t = 1, n + 1,$ $\frac{n}{2} + 2 \leq s \leq n$	$n - 2$
$u_s^t$	7	$4n - 3(s - 1) - t$	$s = 1,$ $2 \leq t \leq \frac{n}{2} + 1$	$\frac{n}{2}$
$u_s^t$	9	$4n - 3(s - 1) - t$	$2 \leq s \leq \frac{n}{2},$ $s + 1 \leq t \leq \frac{n}{2} + 1$	$\frac{n^2}{8} - \frac{3n}{4}$
$u_s^t$	9	$4(n + 1) - s - 3t$	$2 \leq s \leq \frac{n}{2},$ $2 \leq t \leq s$	$\frac{n^2}{8} - \frac{n}{4}$
$u_s^t$	9	$3n + s - 3t + 2$	$\frac{n}{2} + 1 \leq s \leq n - 1,$ $2 \leq t \leq n + 1 - s$	$\frac{n^2}{8} - \frac{n}{4}$
$u_s^t$	9	$n + 3s - t - 1$	$\frac{n}{2} + 1 \leq s \leq n,$ $n - s + 2 \leq t \leq \frac{n}{2} + 1$	$\frac{n^2}{8} + \frac{n}{4}$
$u_s^t$	7	$3(n - s) + t + 1$	$s = 1,$ $\frac{n}{2} + 2 \leq t \leq n$	$\frac{n}{2} - 1$
$u_s^t$	9	$3(n - s) + t + 1$	$2 \leq s \leq \frac{n}{2} - 1,$ $\frac{n}{2} + 2 \leq t \leq n - s + 1$	$\frac{1}{8}(n - 4)(n - 2)$
$u_s^t$	9	$n - s + 3t - 2$	$2 \leq s \leq \frac{n}{2},$ $n - s + 2 \leq t \leq n$	$\frac{n^2}{8} - \frac{n}{4}$
$u_s^t$	9	$4s - n + t - 3$	$\frac{n}{2} + 2 \leq s \leq n,$ $\frac{n}{2} + 2 \leq t \leq s$	$\frac{n^2}{8} - \frac{n}{4}$
$u_s^t$	9	$s + 3t - 4$	$\frac{n}{2} + 1 \leq s \leq n - 1,$ $s + 1 \leq t \leq n$	$\frac{n^2}{8} - \frac{n}{4}$

**Theorem 2.** For every  $n \geq 3$  and  $n \equiv 1 \pmod{2}$  consider the graph of  $G \cong O_n^m$ , with  $n = m$ . Then the modified eccentric connectivity index  $\xi_c(G)$  of  $G$  is equal to

$$\begin{aligned}
\xi_c(O_n^m) &= 225n^2 - 132n + 35 \\
&+ 36 \sum_{s=2}^{\frac{n+1}{2}-1} \left[ \sum_{t=s+1}^{\frac{n+1}{2}} \{4n - 3(s-1) - t\} \right] + 36 \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=2}^s \{4(n+1) - s - 3t\} \right] \\
&+ 36 \sum_{s=\frac{n+1}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} \{3(n-t) + s + 2\} \right] + 36 \sum_{s=\frac{n+1}{2}+1}^n \left[ \sum_{t=n-s+2}^{\frac{n+1}{2}} \{n + 3s - t - 1\} \right] \\
&+ 36 \sum_{s=1}^{\frac{n+1}{2}-1} \left[ \sum_{t=\frac{n+1}{2}+1}^{n-s+1} \{3(n-s) + t + 1\} \right] + 36 \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=n+2-s}^n \{n - s + 3t - 2\} \right] \\
&+ 36 \sum_{s=\frac{n+1}{2}+1}^n \left[ \sum_{t=\frac{n+1}{2}+1}^s \{4s + t - n - 3\} \right] + 36 \sum_{s=\frac{n+1}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n \{s + 3t - 4\} \right].
\end{aligned}$$

*Proof.* Let  $G$  be the graph of  $O_n^m$  and  $n \geq 3$  is odd. As above note that graph of  $O_n^m$  is a symmetric about reflection and rotation at right angles. Thus the eccentricities  $\varepsilon_{u_s^t} = \varepsilon_{v_{n+1-s}^t}$  and from the symmetry at right angles we can obtain that the eccentricities  $\varepsilon_{y_s^t} = \varepsilon_{u_i^s}$ ,  $\varepsilon_{w_i^t} = \varepsilon_{v_i^s}$ . Therefore, by using Table 2 and equation (3) the modified eccentric connectivity index  $\xi_c(G)$ , we get

$$\begin{aligned}
\xi_c(O_n^m) &= 4 \left[ 2 \times 4 \times 4n + 2 \sum_{s=2}^{\frac{n+1}{2}} 5\{4n + 1 - s\} + 2 \sum_{s=\frac{n+1}{2}+1}^n 5\{3n + s - 1\} \right] \\
&+ 4 \left[ \sum_{t=2}^{\frac{n+1}{2}} 7\{4n - t\} + \sum_{s=2}^{\frac{n+1}{2}-1} \left[ \sum_{t=s+1}^{\frac{n+1}{2}} 9\{4n - 3(s-1) - t\} \right] \right] \\
&+ \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=2}^s 9\{4(n+1) - s - 3t\} \right] + \sum_{s=\frac{n+1}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} 9\{3(n-t) + s + 2\} \right] \\
&+ \sum_{s=\frac{n+1}{2}+1}^n \left[ \sum_{t=n-s+2}^{\frac{n+1}{2}} 9\{n + 3s - t - 1\} \right] + \sum_{t=\frac{n+1}{2}+1}^n 7\{3(n-1) + t + 1\} \\
&+ \sum_{s=2}^{\frac{n+1}{2}-1} \left[ \sum_{t=\frac{n+1}{2}+1}^{n-s+1} 9\{3(n-s) + t + 1\} \right] + \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=n+2-s}^n 9\{n - s + 3t - 2\} \right] \\
&+ \sum_{s=\frac{n+1}{2}+1}^n \left[ \sum_{t=\frac{n+1}{2}+1}^s 9\{4s + t - n - 3\} \right] + \sum_{s=\frac{n+1}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n 9\{s + 3t - 4\} \right].
\end{aligned}$$



After some easy calculations we get

$$\begin{aligned}
\xi_c(O_n^m) &= 225n^2 - 132n + 35 \\
&+ 36 \sum_{s=2}^{\frac{n+1}{2}-1} \left[ \sum_{t=s+1}^{\frac{n+1}{2}} \{4n - 3(s-1) - t\} \right] + 36 \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=2}^s \{4(n+1) - s - 3t\} \right] \\
&+ 36 \sum_{s=\frac{n+1}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} \{3(n-t) + s + 2\} \right] + 36 \sum_{s=\frac{n+1}{2}+1}^n \left[ \sum_{t=n-s+2}^{\frac{n+1}{2}} \{n + 3s - t - 1\} \right] \\
&+ 36 \sum_{s=1}^{\frac{n+1}{2}-1} \left[ \sum_{t=\frac{n+1}{2}+1}^{n-s+1} \{3(n-s) + t + 1\} \right] + 36 \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=n+2-s}^n \{n - s + 3t - 2\} \right] \\
&+ 36 \sum_{s=\frac{n+1}{2}+1}^n \left[ \sum_{t=\frac{n+1}{2}+1}^s \{4s + t - n - 3\} \right] + 36 \sum_{s=\frac{n+1}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n \{s + 3t - 4\} \right].
\end{aligned}$$

□

**Theorem 3.** For every  $n \geq 4$  and  $n \equiv 0 \pmod{2}$  consider the graph of  $G \cong O_n^m$ , with  $n = m$ . Then the Ediz eccentric connectivity index of  $G$  is equal to

$$\begin{aligned}
E_{\xi^c}(O_n^m) &= \frac{8}{n} + 40 \sum_{s=2}^{\frac{n}{2}+1} \frac{1}{4n+1-s} + 40 \sum_{s=\frac{n}{2}+2}^n \frac{1}{3n+s-1} \\
&+ 28 \sum_{t=2}^{\frac{n}{2}+1} \frac{1}{4n-t} + 28 \sum_{t=\frac{n}{2}+2}^n \frac{1}{3(n-1)+t+1} \\
&+ 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} \frac{1}{4n-3(s-1)-t} \right] + 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s \frac{1}{4(n+1)-s-3t} \right] \\
&+ 36 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} \frac{1}{3(n-t)+s+2} \right] + 36 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} \frac{1}{n+3s-t-1} \right] \\
&+ 36 \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} \frac{1}{3(n-s)+t+1} \right] + 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n \frac{1}{n-s+3t-2} \right] \\
&+ 36 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s \frac{1}{4s+t-n-3} \right] + 36 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n \frac{1}{s+3t-4} \right].
\end{aligned}$$

TABLE 2. Partition of vertices of the type  $u_s^t$  of  $O_n^m$  based on degree sum and eccentricity of each vertex when  $n \equiv 1 \pmod{2}$ .

Representative	$S_{u_s^t}$	eccentricity	Range	Frequency
$u_s^t$	4	$4n - s + 1$	$t = 1, n + 1; s = 1$	2
$u_s^t$	5	$4n - s + 1$	$t = 1, n + 1,$ $2 \leq s \leq \frac{n+1}{2}$	$n - 1$
$u_s^t$	5	$3n + s - 1$	$t = 1, n + 1,$ $\frac{n+1}{2} + 1 \leq s \leq n$	$n - 1$
$u_s^t$	7	$4n - 3(s - 1) - t$	$1 = s,$ $2 \leq t \leq \frac{n+1}{2}$	$(\frac{n+1}{2} - 1)$
$u_s^t$	9	$4n - 3(s - 1) - t$	$2 \leq s \leq \frac{n+1}{2} - 1,$ $s + 1 \leq t \leq \frac{n+1}{2}$	$\frac{n-3}{4}(\frac{n+1}{2} - 1)$
$u_s^t$	9	$4(n + 1) - s - 3t$	$2 \leq s \leq \frac{n+1}{2},$ $2 \leq t \leq s$	$\frac{n+1}{4}(\frac{n+1}{2} - 1)$
$u_s^t$	9	$3n + s - 3t + 2$	$\frac{n+1}{2} + 1 \leq s \leq n - 1,$ $2 \leq t \leq n + 1 - s$	$\frac{n-1}{4}(\frac{n-1}{2} - 1)$
$u_s^t$	9	$n + 3s - t - 1$	$\frac{n+1}{2} + 1 \leq s \leq n,$ $n - s + 2 \leq t \leq \frac{n+1}{2}$	$\frac{n-1}{4}(\frac{n-1}{2} + 1)$
$u_s^t$	7	$3(n - s) + t + 1$	$s = 1,$ $\frac{n+1}{2} + 1 \leq t \leq n$	$(\frac{n+1}{2} - 1)$
$u_s^t$	9	$n - s + 3t - 2$	$2 \leq s \leq \frac{n+1}{2},$ $n - s + 2 \leq t \leq n$	$\frac{n-3}{4}(\frac{n+1}{2} - 1)$
$u_s^t$	9	$4s - n + t - 3$	$\frac{n+1}{2} + 1 \leq s \leq n,$ $\frac{n+1}{2} + 1 \leq t \leq s$	$\frac{n-1}{4}(\frac{n+1}{2})$
$u_s^t$	9	$s + 3t - 4$	$\frac{n+1}{2} + 1 \leq s \leq n - 1,$ $s + 1 \leq t \leq n$	$\frac{n-1}{4}(\frac{n-1}{2} - 1)$

*Proof.* Let  $G$  be the graph of  $O_n^m$  and  $n \equiv 0 \pmod{2}$ . By using the arguments in proof of Theorem 1, Table 1 and following formula the Ediz eccentric connectivity index  ${}^E\xi^c(G)$  of  $O_n^m$  is equal to

$${}^E\xi^c(G) = \sum_{v \in V(G)} \left( \frac{S_v}{\epsilon_v} \right) = 4 \sum_{u_s^t \in V(G)} \left( \frac{S_{u_s^t}}{\epsilon_{u_s^t}} \right)$$

$$\begin{aligned}
E_{\xi^c}(O_n^m) &= 4 \left[ 2 \times \frac{4}{4n} + 2 \sum_{s=2}^{\frac{n}{2}+1} \frac{5}{4n+1-s} + 2 \sum_{s=\frac{n}{2}+2}^n \frac{5}{3n+s-1} \right] \\
&+ 4 \left[ \sum_{t=2}^{\frac{n}{2}+1} \frac{7}{4n-t} + \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} \frac{9}{4n-3(s-1)-t} \right] \right] \\
&+ \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s \frac{9}{4(n+1)-s-3t} \right] + \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} \frac{9}{3(n-t)+s+2} \right] \\
&+ \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} \frac{9}{n+3s-t-1} \right] + \sum_{t=\frac{n}{2}+2}^n \frac{7}{3(n-1)+t+1} \\
&+ \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} \frac{9}{3(n-s)+t+1} \right] + \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n \frac{9}{n-s+3t-2} \right] \\
&+ \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s \frac{9}{4s+t-n-3} \right] + \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n \frac{9}{s+3t-4} \right].
\end{aligned}$$

After an easy computation, we get

$$\begin{aligned}
E_{\xi^c}(O_n^m) &= \frac{8}{n} + 40 \sum_{s=2}^{\frac{n}{2}+1} \frac{1}{4n+1-s} + 40 \sum_{s=\frac{n}{2}+2}^n \frac{1}{3n+s-1} \\
&+ 28 \sum_{t=2}^{\frac{n}{2}+1} \frac{1}{4n-t} + 28 \sum_{t=\frac{n}{2}+2}^n \frac{1}{3(n-1)+t+1} \\
&+ 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} \frac{1}{4n-3(s-1)-t} \right] + 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s \frac{1}{4(n+1)-s-3t} \right] \\
&+ 36 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} \frac{1}{3(n-t)+s+2} \right] + 36 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} \frac{1}{n+3s-t-1} \right] \\
&+ 36 \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} \frac{1}{3(n-s)+t+1} \right] + 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n \frac{1}{n-s+3t-2} \right] \\
&+ 36 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s \frac{1}{4s+t-n-3} \right] + 36 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n \frac{1}{s+3t-4} \right].
\end{aligned}$$

□

**Theorem 4.** For every  $n \geq 3$  and  $n \equiv 1 \pmod{2}$  consider the graph of  $G \cong O_n^m$ , with  $n = m$ . Then the Ediz eccentric connectivity index of  $G$  is equal to

$$\begin{aligned}
E_{\xi^c}(O_n^m) &= \frac{8}{n} + 40 \sum_{s=2}^{\frac{n+1}{2}} \frac{1}{4n+1-s} + 40 \sum_{s=\frac{\frac{n+1}{2}+1}^n} \frac{1}{3n+s-1} \\
&+ 28 \sum_{t=2}^{\frac{n+1}{2}} \frac{1}{4n-t} + 28 \sum_{t=\frac{\frac{n+1}{2}+1}^n} \frac{1}{3(n-1)+t+1} \\
&+ 36 \sum_{s=2}^{\frac{n+1}{2}-1} \left[ \sum_{t=s+1}^{\frac{n+1}{2}} \frac{1}{4n-3(s-1)-t} \right] + 36 \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=2}^s \frac{1}{4(n+1)-s-3t} \right] \\
&+ 36 \sum_{s=\frac{\frac{n+1}{2}+1}^{n-1}} \left[ \sum_{t=2}^{n-s+1} \frac{1}{3(n-t)+s+2} \right] + 36 \sum_{s=\frac{\frac{n+1}{2}+1}^n} \left[ \sum_{t=n-s+2}^{\frac{n+1}{2}} \frac{1}{n+3s-t-1} \right] \\
&+ 36 \sum_{s=2}^{\frac{n+1}{2}-1} \left[ \sum_{t=\frac{\frac{n+1}{2}+1}^n} \frac{1}{3(n-s)+t+1} \right] + 36 \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=n+2-s}^n \frac{1}{n-s+3t-2} \right] \\
&+ 36 \sum_{s=\frac{\frac{n+1}{2}+1}^n} \left[ \sum_{t=\frac{\frac{n+1}{2}+1}^s} \frac{1}{4s+t-n-3} \right] + 36 \sum_{s=\frac{\frac{n+1}{2}+1}^{n-1}} \left[ \sum_{t=s+1}^n \frac{1}{s+3t-4} \right].
\end{aligned}$$

*Proof.* Let  $G$  be the graph of  $O_n^m$  and  $n \equiv 1 \pmod{2}$ . By using the arguments in proof of Theorem 2, Table 2 and from formula (4) the result follows.  $\square$

**Theorem 5.** For every  $n \geq 4$  and  $n \equiv 0 \pmod{2}$  consider the graph of  $G \cong O_n^m$ , with  $n = m$ . Then the modified eccentric connectivity polynomial of  $G$  is equal to

$$\begin{aligned}
\xi_c(O_n^m, x) &= \frac{1}{x-1} \left( (-28x^{4n-1} + 40x^{3n} - 40x^{4n}) \left(\frac{1}{x}\right)^{\binom{n}{2}} - 40x^{(3n+1)} \left(\frac{1}{x}\right)^n + 24x^{4n} \right. \\
&\quad \left. - 28x^{(7n/2)} + 56x^{4n-1} + 16x^{4n+1} \right) \\
&+ 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} x^{(4n-3(s-1)-t)} \right] + 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s x^{(4(n+1)-s-3t)} \right] \\
&+ 36 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} x^{(3(n-t)+s+2)} \right] + 36 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} x^{(n+3s-t-1)} \right] \\
&+ 36 \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} x^{(3(n-s)+t+1)} \right] + 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n x^{(n-s+3t-2)} \right] \\
&+ 36 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s x^{(4s+t-n-3)} \right] + 36 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n x^{(s+3t-4)} \right].
\end{aligned}$$

*Proof.* By using the arguments in the proof of Theorem 1, the values from Table 1 and equation (5) given below we get

$$\xi_c(G, x) = \sum_{u \in V(G)} S_u x^{\epsilon_u} = 4 \sum_{u_s^t \in V(G)} S_{u_s^t} x^{\epsilon_{u_s^t}}$$

$$\begin{aligned}
\xi_c(O_n^m, x) &= 4 \left[ 2 \times 4x^{4n} + 2 \sum_{s=2}^{\frac{n}{2}+1} 5x^{(4n+1-s)} + 2 \sum_{s=\frac{n}{2}+2}^n 5x^{(3n+s-1)} \right] \\
&+ 4 \left[ \sum_{t=2}^{\frac{n}{2}+1} 7x^{(4n-t)} + \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} 9x^{(4n-t)} \right] \right] \\
&+ \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s 9x^{(4(n+1)-s-3t)} \right] + \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} 9x^{(3(n-t)+s+2)} \right] \\
&+ \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} 9x^{(n+3s-t-1)} \right] + \sum_{t=\frac{n}{2}+2}^n 7x^{(3(n-1)+t+1)} \\
&+ \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} 9x^{(3(n-s)+t+1)} \right] + \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n 9x^{(n-s+3t-2)} \right] \\
&+ \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s 9x^{(4s+t-n-3)} \right] + \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n 9x^{(s+3t-4)} \right].
\end{aligned}$$

After some easy calculations we get

$$\begin{aligned}
\xi_c(O_n^m, x) &= \frac{1}{x-1} \left( (-28x^{4n-1} + 40x^{3n} - 40x^{4n}) \left( \frac{1}{x} \right)^{\binom{n}{2}} - 40x^{(3n+1)} \left( \frac{1}{x} \right)^n + 24x^{4n} \right. \\
&- \left. 28x^{(7n/2)} + 56x^{4n-1} + 16x^{4n+1} \right) \\
&+ 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} x^{(4n-3(s-1)-t)} \right] + 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s x^{(4(n+1)-s-3t)} \right] \\
&+ 36 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} x^{(3(n-t)+s+2)} \right] + 36 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} x^{(n+3s-t-1)} \right] \\
&+ 36 \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} x^{(3(n-s)+t+1)} \right] + 36 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n x^{(n-s+3t-2)} \right] \\
&+ 36 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s x^{(4s+t-n-3)} \right] + 36 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n x^{(s+3t-4)} \right].
\end{aligned}$$

□

**Theorem 6.** For every  $n \geq 3$  and  $n \equiv 1 \pmod{2}$  consider the graph of  $G \cong O_n^m$ , with  $n = m$ . Then the modified eccentric connectivity polynomial of  $G$  is equal to

$$\begin{aligned} \xi_c(O_n^m, x) &= \frac{1}{x-1} \left( (40x^{3n+2} - 28x^{4n+1} - 40x^{4n+2}) \left( \frac{1}{x} \right)^{\left( \frac{n}{2} + \frac{3}{2} \right)} - 40x^{(3n+1)} \left( \frac{1}{x} \right)^n + 24x^{4n} \right. \\ &\quad \left. - 28x^{\left( \frac{7n}{2} \right) - \frac{1}{2}} + 56x^{4n-1} + 16x^{4n+1} \right) \\ &+ 36 \sum_{s=2}^{\frac{n+1}{2}-1} \left[ \sum_{t=s+1}^{\frac{n+1}{2}} x^{(4n-3(s-1)-t)} \right] + 36 \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=2}^s x^{(4(n+1)-s-3t)} \right] \\ &+ 36 \sum_{s=\frac{n+1}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} x^{(3(n-t)+s+2)} \right] + 36 \sum_{s=\frac{n+1}{2}+1}^n \left[ \sum_{t=n-s+2}^{\frac{n+1}{2}} x^{(n+3s-t-1)} \right] \\ &+ 36 \sum_{s=1}^{\frac{n+1}{2}-1} \left[ \sum_{t=\frac{n+1}{2}+1}^{n-s+1} x^{(3(n-s)+t+1)} \right] + 36 \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=n+2-s}^n x^{(n-s+3t-2)} \right] \\ &+ \sum_{s=\frac{n+1}{2}+1}^n \left[ \sum_{t=\frac{n+1}{2}+1}^s x^{(4s+t-n-3)} \right] + 36 \sum_{s=\frac{n+1}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n x^{(s+3t-4)} \right]. \end{aligned}$$

*Proof.* Let  $G \cong O_n^m$ ,  $n \geq 3$  and  $n \equiv 1 \pmod{2}$ . By using the arguments in the proof of Theorem 1, as in Theorem 5, the values from Table 2 and equation (5) the result follows.  $\square$

In Table 3 and Table 4 we have partitioned the vertices of the type  $u_s^t$  of  $O_n^m$  based on degree product and eccentricity of each vertex. This will help us to develop the coming theorems.

**Theorem 7.** For every  $n \geq 4$  and  $n \equiv 0 \pmod{2}$  consider the graph of  $G \cong O_n^m$ , with  $n = m$ . Then the modified augmented eccentric connectivity index  $^{MA}\xi_c^c(G)$  of  $G$  is equal to

$$\begin{aligned}
MA_{\xi_c^c}(O_n^m) &= 324n^2 - 232n + 48 \\
&+ 108 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} \{4n - 3(s-1) - t\} \right] + 108 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s \{4(n+1) - s - 3t\} \right] \\
&+ 108 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} \{3(n-t) + s + 2\} \right] + 108 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} \{n + 3s - t - 1\} \right] \\
&+ 108 \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} \{3(n-s) + t + 1\} \right] + 108 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n \{n - s + 3t - 2\} \right] \\
&+ 108 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s \{4s + t - n - 3\} \right] + 108 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n \{s + 3t - 4\} \right].
\end{aligned}$$

*Proof.* Let  $G$  be the graph of  $O_n^m$ . Therefore, from Table 3 and formula (8), given below, the modified augmented eccentric connectivity index  $MA_{\xi_c^c}(O_n^m)$  of  $O_n^m$  can be calculated. Hence the result.

$$(8) \quad MA_{\xi_c^c}(G, x) = \sum_{v \in V(G)} M_v \epsilon_v$$

$$MA_{\xi_c^c}(O_n^m) = 4 \sum_{u_s^t \in V(G)} (M_{u_s^t} \epsilon_{u_s^t})$$

$$\begin{aligned}
MA_{\xi_c^c}(O_n^m) &= 4 \left[ 2 \times 4 \times 4n + 2 \sum_{s=2}^{\frac{n}{2}+1} 6\{4n + 1 - s\} + 2 \sum_{s=\frac{n}{2}+2}^n 6\{3n + s - 1\} \right] \\
&+ 4 \left[ \sum_{t=2}^{\frac{n}{2}+1} 12(4n - t) + \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} 27\{4n - 3(s-1) - t\} \right] \right]
\end{aligned}$$



$$\begin{aligned}
& + \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s 27\{4(n+1) - s - 3t\} \right] + \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} 27\{3(n-t) + s + 2\} \right] \\
& + \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} 27\{n + 3s - t - 1\} \right] + \sum_{t=\frac{n}{2}+2}^n 12\{3(n-1) + t + 1\} \\
& + \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} 27\{3(n-s) + t + 1\} \right] + \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n 27\{n-s+3t-2\} \right] \\
& + \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s 27\{4s+t-n-3\} \right] + \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n 27\{s+3t-4\} \right].
\end{aligned}$$

After some easy calculations we get

$$\begin{aligned}
MA_{\xi_c}(O_n^m) & = 324n^2 - 232n + 48 \\
& + 108 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} \{4n - 3(s-1) - t\} \right] + 108 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s \{4(n+1) - s - 3t\} \right] \\
& + 108 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} \{3(n-t) + s + 2\} \right] + 108 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} \{n + 3s - t - 1\} \right] \\
& + 108 \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} \{3(n-s) + t + 1\} \right] + 108 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n \{n-s+3t-2\} \right] \\
& + 108 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s \{4s+t-n-3\} \right] + 108 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n \{s+3t-4\} \right].
\end{aligned}$$

□

**Theorem 8.** For every  $n \geq 3$  and  $n \equiv 1 \pmod{2}$  consider the graph of  $G \cong O_n^m$ , with  $n = m$ . Then the modified augmented eccentric connectivity index  $\xi_c(G)$  of  $G$  is equal

to

$$\begin{aligned}
MA_{\xi_c^c}(O_n^m) &= 324n^2 - 256n + 60 \\
&+ 108 \sum_{s=2}^{\frac{n+1}{2}-1} \left[ \sum_{t=s+1}^{\frac{n+1}{2}} \{4n - 3(s-1) - t\} \right] + 108 \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=2}^s \{4(n+1) - s - 3t\} \right] \\
&+ 108 \sum_{s=\frac{n+1}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} \{3(n-t) + s + 2\} \right] + 108 \sum_{s=\frac{n+1}{2}+1}^n \left[ \sum_{t=n-s+2}^{\frac{n+1}{2}} \{n + 3s - t - 1\} \right] \\
&+ 108 \sum_{s=1}^{\frac{n+1}{2}-1} \left[ \sum_{t=\frac{n+1}{2}+1}^{n-s+1} \{3(n-s) + t + 1\} \right] + 108 \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=n+2-s}^n \{n - s + 3t - 2\} \right] \\
&+ 108 \sum_{s=\frac{n+1}{2}+1}^n \left[ \sum_{t=\frac{n+1}{2}+1}^s \{4s + t - n - 3\} \right] + 108 \sum_{s=\frac{n+1}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n \{s + 3t - 4\} \right].
\end{aligned}$$

*Proof.* As in Theorem 7, by using Table 4 and equation (8) the result follows.  $\square$

**Theorem 9.** For every  $n \geq 4$  and  $n \equiv 0 \pmod{2}$  consider the graph of  $G \cong O_n^m$ , with  $n = m$ . Then the modified augmented eccentric connectivity polynomial of  $G$  is equal to

$$\begin{aligned}
MA_{\xi_c^c}(O_n^m, x) &= \frac{1}{x-1} \left( (48x^{4n-1} + 48x^{3n} - 48x^{4n}) \left( \frac{1}{x} \right)^{\binom{n}{2}} - 48x^{(3n+1)} \left( \frac{1}{x} \right)^n + 32x^{4n} \right. \\
&\quad \left. - 48x^{(7n/2)} + 96x^{4n-1} + 16x^{4n+1} \right) \\
&+ 48 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} x^{(4n-3(s-1)-t)} \right] + 48 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s x^{(4(n+1)-s-3t)} \right] \\
&+ 48 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} x^{(3(n-t)+s+2)} \right] + 48 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} x^{(n+3s-t-1)} \right] \\
&+ 48 \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} x^{(3(n-s)+t+1)} \right] + 48 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n x^{(n-s+3t-2)} \right] \\
&+ 48 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s x^{(4s+t-n-3)} \right] + 48 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n x^{(s+3t-4)} \right].
\end{aligned}$$

TABLE 3. Partition of vertices of the type  $u_s^t$  of  $O_n^m$  based on degree product and eccentricity of each vertex when  $n \equiv 0 \pmod{2}$ .

Representative	$M_{u_s^t}$	eccentricity	Range	Frequency
$u_s^t$	4	$4n - s + 1$	$t = 1, n + 1; s = 1$	2
$u_s^t$	6	$4n - s + 1$	$t = 1, n + 1,$ $2 \leq s \leq \frac{n}{2} + 1$	$n$
$u_s^t$	6	$3n + s - 1$	$t = 1, n + 1,$ $\frac{n}{2} + 2 \leq s \leq n$	$n - 2$
$u_s^t$	12	$4n - 3(s - 1) - t$	$s = 1,$ $2 \leq t \leq \frac{n}{2} + 1$	$\frac{n}{2}$
$u_s^t$	27	$4n - 3(s - 1) - t$	$2 \leq s \leq \frac{n}{2},$ $s + 1 \leq t \leq \frac{n}{2} + 1$	$\frac{n^2}{8} - \frac{3n}{4}$
$u_s^t$	27	$4(n + 1) - s - 3t$	$2 \leq s \leq \frac{n}{2},$ $2 \leq t \leq s$	$\frac{n}{4}(\frac{n}{2} - 1)$
$u_s^t$	27	$3n + s - 3t + 2$	$\frac{n}{2} + 1 \leq s \leq n - 1,$ $2 \leq t \leq n + 1 - s$	$\frac{n}{4}(\frac{n}{2} - 1)$
$u_s^t$	27	$n + 3s - t - 1$	$\frac{n}{2} + 1 \leq s \leq n,$ $n - s + 2 \leq t \leq \frac{n}{2} + 1$	$\frac{n}{4}(\frac{n}{2} + 1)$
$u_s^t$	12	$3(n - s) + t + 1$	$s = 1,$ $\frac{n}{2} + 2 \leq t \leq n$	$\frac{n}{2} - 1$
$u_s^t$	27	$3(n - s) + t + 1$	$2 \leq s \leq \frac{n}{2} - 1,$ $\frac{n}{2} + 2 \leq t \leq n - s + 1$	$\frac{1}{8}(n - 4)(n - 2)$
$u_s^t$	27	$n - s + 3t - 2$	$2 \leq s \leq \frac{n}{2},$ $n - s + 2 \leq t \leq n$	$\frac{n}{4}(\frac{n}{2} - 1)$
$u_s^t$	27	$4s - n + t - 3$	$\frac{n}{2} + 2 \leq s \leq n,$ $\frac{n}{2} + 2 \leq t \leq s$	$\frac{n}{4}(\frac{n}{2} - 1)$
$u_s^t$	27	$s + 3t - 4$	$\frac{n}{2} + 1 \leq s \leq n - 1,$ $s + 1 \leq t \leq n$	$\frac{n}{4}(\frac{n}{2} - 1)$

*Proof.* By using the arguments in the proof of Theorem 1, the values from Table 3 and equation (8) given below we get

$${}^{MA}\xi_c^c(G, x) = \sum_{u \in V(G)} M_u x^{\epsilon_u} = 4 \sum_{u_s^t \in V(G)} M_{u_s^t} x^{\epsilon_{u_s^t}}$$

TABLE 4. Partition of vertices of the type  $u_s^t$  of  $O_n^m$  based on degree product and eccentricity of each vertex when  $n \equiv 1 \pmod{2}$ .

Representative	$M_{u_s^t}$	eccentricity	Range	Frequency
$u_s^t$	4	$4n - s + 1$	$t = 1, n + 1; s = 1$	2
$u_s^t$	6	$4n - s + 1$	$t = 1, n + 1,$ $2 \leq s \leq \frac{n+1}{2}$	$n - 1$
$u_s^t$	6	$3n + s - 1$	$t = 1, n + 1,$ $\frac{n+1}{2} + 1 \leq s \leq n$	$n - 1$
$u_s^t$	12	$4n - 3(s - 1) - t$	$1 = s,$ $2 \leq t \leq \frac{n+1}{2}$	$(\frac{n+1}{2} - 1)$
$u_s^t$	27	$4n - 3(s - 1) - t$	$2 \leq s \leq \frac{n+1}{2} - 1,$ $s + 1 \leq t \leq \frac{n+1}{2}$	$\frac{n-3}{4}(\frac{n+1}{2} - 1)$
$u_s^t$	27	$4(n + 1) - s - 3t$	$2 \leq s \leq \frac{n+1}{2},$ $2 \leq t \leq s$	$\frac{n+1}{4}(\frac{n+1}{2} - 1)$
$u_s^t$	27	$3n + s - 3t + 2$	$\frac{n+1}{2} + 1 \leq s \leq n - 1,$ $2 \leq t \leq n + 1 - s$	$\frac{n-1}{4}(\frac{n-1}{2} - 1)$
$u_s^t$	27	$n + 3s - t - 1$	$\frac{n+1}{2} + 1 \leq s \leq n,$ $n - s + 2 \leq t \leq \frac{n+1}{2}$	$\frac{n-1}{4}(\frac{n-1}{2} + 1)$
$u_s^t$	12	$3(n - s) + t + 1$	$s = 1,$ $\frac{n+1}{2} + 1 \leq t \leq n$	$(\frac{n+1}{2} - 1)$
$u_s^t$	27	$n - s + 3t - 2$	$2 \leq s \leq \frac{n+1}{2},$ $n - s + 2 \leq t \leq n$	$\frac{n-3}{4}(\frac{n+1}{2} - 1)$
$u_s^t$	27	$4s - n + t - 3$	$\frac{n+1}{2} + 1 \leq s \leq n,$ $\frac{n+1}{2} + 1 \leq t \leq s$	$\frac{n-1}{4}(\frac{n+1}{2})$
$u_s^t$	27	$s + 3t - 4$	$\frac{n+1}{2} + 1 \leq s \leq n - 1,$ $s + 1 \leq t \leq n$	$\frac{n-1}{4}(\frac{n-1}{2} - 1)$

$$\begin{aligned}
MA\xi_c(O_n^m, x) &= 4 \left[ 2 \times 4x^{4n} + 2 \sum_{s=2}^{\frac{n}{2}+1} 6x^{(4n+1-s)} + 2 \sum_{s=\frac{n}{2}+2}^n 6x^{(3n+s-1)} \right] \\
&+ 4 \left[ \sum_{t=2}^{\frac{n}{2}+1} 12x^{(4n-t)} + \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} 27x^{(4n-t)} \right] \right]
\end{aligned}$$

$$\begin{aligned}
& + \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s 27x^{(4(n+1)-s-3t)} \right] + \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} 27x^{(3(n-t)+s+2)} \right] \\
& + \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} 27x^{(n+3s-t-1)} \right] + \sum_{t=\frac{n}{2}+2}^n 12x^{(3(n-1)+t+1)} \\
& + \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} 27x^{(3(n-s)+t+1)} \right] + \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n 27x^{(n-s+3t-2)} \right] \\
& + \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s 27x^{(4s+t-n-3)} \right] + \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n 27x^{(s+3t-4)} \right].
\end{aligned}$$

After some easy calculations we get

$$\begin{aligned}
MA\xi_c^c(O_n^m, x) &= \frac{1}{x-1} \left( (48x^{4n-1} + 48x^{3n} - 48x^{4n}) \left( \frac{1}{x} \right)^{\binom{n}{2}} - 48x^{(3n+1)} \left( \frac{1}{x} \right)^n + 32x^{4n} \right. \\
& \left. - 48x^{(7n/2)} + 96x^{4n-1} + 16x^{4n+1} \right) \\
& + 48 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=s+1}^{\frac{n}{2}+1} x^{(4n-3(s-1)-t)} \right] + 48 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=2}^s x^{(4(n+1)-s-3t)} \right] \\
& + 48 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} x^{(3(n-t)+s+2)} \right] + 48 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=n-s+2}^{\frac{n}{2}+1} x^{(n+3s-t-1)} \right] \\
& + 48 \sum_{s=2}^{\frac{n}{2}-1} \left[ \sum_{t=\frac{n}{2}+2}^{n-s+1} x^{(3(n-s)+t+1)} \right] + 48 \sum_{s=2}^{\frac{n}{2}} \left[ \sum_{t=n+2-s}^n x^{(n-s+3t-2)} \right] \\
& + 48 \sum_{s=\frac{n}{2}+2}^n \left[ \sum_{t=\frac{n}{2}+2}^s x^{(4s+t-n-3)} \right] + 48 \sum_{s=\frac{n}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n x^{(s+3t-4)} \right].
\end{aligned}$$

□

**Theorem 10.** For every  $n \geq 3$  and  $n \equiv 1 \pmod{2}$  consider the graph of  $G \cong O_n^m$ , with  $n = m$ . Then the modified augmented eccentric connectivity polynomial of  $G$  is equal to

$$\begin{aligned}
MA\xi_c^c(O_n^m, x) &= \frac{1}{x-1} \left( (-48x^{3n+2} - 48x^{4n+1} - 48x^{4n+2}) \left(\frac{1}{x}\right)^{\left(\frac{n}{2} + \frac{3}{2}\right)} - 48x^{(3n+1)} \left(\frac{1}{x}\right)^n + 32x^{4n} \right. \\
&\quad \left. - 48x^{\left(\frac{7n}{2}\right) - \frac{1}{2}} + 96x^{4n-1} + 16x^{4n+1} \right) \\
&+ 108 \sum_{s=2}^{\frac{n+1}{2}-1} \left[ \sum_{t=s+1}^{\frac{n+1}{2}} x^{(4n-3(s-1)-t)} \right] + 108 \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=2}^s x^{(4(n+1)-s-3t)} \right] \\
&+ 108 \sum_{s=\frac{n+1}{2}+1}^{n-1} \left[ \sum_{t=2}^{n-s+1} x^{(3(n-t)+s+2)} \right] + 108 \sum_{s=\frac{n+1}{2}+1}^n \left[ \sum_{t=n-s+2}^{\frac{n+1}{2}} x^{(n+3s-t-1)} \right] \\
&+ 108 \sum_{s=1}^{\frac{n+1}{2}-1} \left[ \sum_{t=\frac{n+1}{2}+1}^{n-s+1} x^{(3(n-s)+t+1)} \right] + 108 \sum_{s=2}^{\frac{n+1}{2}} \left[ \sum_{t=n+2-s}^n x^{(n-s+3t-2)} \right] \\
&+ \sum_{s=\frac{n+1}{2}+1}^n \left[ \sum_{t=\frac{n+1}{2}+1}^s x^{(4s+t-n-3)} \right] + 108 \sum_{s=\frac{n+1}{2}+1}^{n-1} \left[ \sum_{t=s+1}^n x^{(s+3t-4)} \right].
\end{aligned}$$

*Proof.* As in Theorem 9, by using Table 4 and the equation (8) the result follows.  $\square$

#### 4. CONCLUSIONS AND COMPARISON BETWEEN THE INDICES

High discriminating power and extremely low degeneracy are desirable properties of an ideal topological index, which researchers in theoretical chemistry are striving to achieve. The values of  $MA\xi^c(G)$  were computed for all the possible structure of three and four vertices. The values and the structures have been presented in Table 5 and their comparison is presented in Table 6. Modified augmented eccentric connectivity index demonstrate exceptionally high discriminating power, defined as the ratio of the highest to lowest value for all possible structures with the same number of vertices. This is evident from the fact that the ratio of the highest to lowest value for all possible structure containing three and four vertices is very high in contrast to  $\xi(G)$  and  $\xi_c(G)$ . The ratio of the highest to lowest value for all possible structures containing four vertices for  $MA\xi^c(G)$  is 6.75 in comparison to 1.78 and 1.7 for  $\xi(G)$  and  $\xi_c(G)$ , respectively. The exceptionally high discriminating power of the proposed indices makes them extremely sensitive towards minor change(s) in molecular structure. This extreme sensitivity towards branching and the discriminating power of proposed indices are clearly evident from the respective index values of all the possible structures with four vertices. Degeneracy: the number of compounds having identical values/the total number of compounds with the same number of vertices.

TABLE 5. Values of eccentric connectivity index, modified eccentric connectivity index and modified augmented eccentric connectivity index for all possible structures with three and four.









S.N	Structure	$\xi(G)$	$^{MA}\xi^c(G)$	$\xi_c(G)$
1		6	9	10
2		6	12	12
3		14	16	24
4		9	19	21
5		13	32	32
6		16	32	32
7		14	60	29
8		12	108	36

TABLE 6. Comparison of the discriminating power and degeneracy of eccentric connectivity index, modified eccentric connectivity index and modified augmented eccentric connectivity index using all possible structures with three and four vertices.

	$\xi(G)$	$^{MA}\xi^c(G)$	$\xi_c(G)$
• For three vertices			
Minimum value	6	9	10
Maximum value	6	12	12
Ratio	1:1	1:1.34	1:1.2
Degeneracy	1/2	0/2	0/2
• For four vertices			
Minimum value	9	16	21
Maximum value	16	108	36
Ratio	1:1.78	1:6.75	1:1.7
Degeneracy	1/6	1/6	1/6

Degeneracy is a measure of the ability of an index to differentiate between the relative positions of atom in a molecule.  $^{MA}\xi^c(G)$  did not exhibit any degeneracy for all possible structures with three vertices whereas  $^{MA}\xi^c(G)$  had a very low degeneracy of one in the

case of all possible structures with four vertices (Table 6).  $\xi(G)$  had one identical values out of 6 structures with only four vertices. Extremely low degeneracy indicates the enhanced capability of these indices to differentiate and demonstrate slight variations in the molecular structure, which clearly reveals the remote chance of different structures having the same value.

The Table 7 shows a comparison between the eccentric connectivity index, modified eccentric connectivity index and modified augmented eccentric connectivity index for octagonal grid  $O_m^n$  for finite  $n = 3, \dots, 10$ .

TABLE 7. comparison of  $\xi_c(O_m^n)$ ,  ${}^E\xi^c(O_m^n)$  and  ${}^{MA}\xi^c(O_m^n)$  for  $O_m^n$ , when  $m = n$ .

$[n, m]$	$\xi_c(O_m^n)$	${}^E\xi^c(O_m^n)$	${}^{MA}\xi^c(O_m^n)$
[3,3]	2888	$\frac{5369}{165}$	5880
[4,4]	6564	$\frac{616039}{15015}$	14456
[5,5]	13460	$\frac{74609462}{1322685}$	32260
[6,6]	22972	$\frac{563513878}{8580495}$	56868
[7,7]	37280	$\frac{803471620793}{10039179150}$	95576
[8,8]	55364	$\frac{1361165885969}{15168440430}$	144424
[9,9]	79784	$\frac{1929246726361}{18627909300}$	212136
[10,10]	109176	$\frac{1149037176620287}{10119188365650}$	293432

#### ACKNOWLEDGEMENTS

The third author of this work were partially supported by MINECO grant number MTM2014-51891-P, Fundación Séneca de la Región de Murcia grant number 19219/PI/14. The third and four authors of this work were partially supported by National Nature Science Foundation of China (Western Region Funds) grant number 11761083.



## REFERENCES

- [1] A. R. Ashrafi and M. Ghorbani, *A study of fullerenes by MEC polynomials*. Electronic Materials Letters, , **vol. 6, no. 2** (2010), pp. 87–90.
- [2] J. Asadpour and L. Safikhani, *Study of  $CNC_7[n]$  Carbon Nanocone by M-eccentric Connectivity Polynomial*. Australian Journal of Basic and Applied Sciences, **7** (2013), 883.
- [3] B. Basavanagoud, V. R. Desai, and S. Patil,  $(\beta, \alpha)$ -Connectivity index of graphs, Applied Mathematics and Nonlinear Sciences, vol. 2, no. 1, pp. 21-30, 2017.
- [4] M. V. Diudea, *Distance counting in tubes and tori: Wiener index and Hosoya polynomial*. In: Nanostructures-NovelArchitecture, NOVA: New York, (2005); pp 203–242.
- [5] T. Došlić, M. Saheli: *Eccentric connectivity index of composite graphs*. Util. Math. **95**(2014) , 3–22.
- [6] N. De, A. Pal, S. M. A. Nayeem, *Modified eccentric Connectivity of Generalized Thorn Graphs*. International Journal of Computational Mathematics (2014), Article ID 436140, 1-8.
- [7] N. De, S. M. A. Nayeem and A. Pal, *Total eccentricity index of the generalized index and polynomial of thorn graph*. Applied Mathematics, **3** (2012), 931-934.
- [8] S. Ediz, *Computing Ediz eccentric connectivity index of an infinite class of nanostar dendrimers*. Optoelectron. Adv. Mater. Rapid Commun. **4** (2010), 1847-1848.
- [9] M. R. Farahani, *Eccentricity Version of Atom-Bond Connectivity Index of Benzenoid Family  $ABC_5(H_k)$* . World Applied Sciences Journal, **21(9)** (2013), 1260–1265.
- [10] Y. Guihai, Q. Hui, L. Tang, L. Feng, *On the connective eccentricity index of trees and unicyclic graphs with given diameter*. Journal of Mathematical Analysis and Applications, **420** (2014, ), 1776–1786.
- [11] S. Gupta, M. Singh, and A. K. Madan, *Connective eccentricity index: a novel topological descriptor for predicting biological activity*. J. Mol. Graph. Model. **18** (2000), 18–25.
- [12] S. Gupta, M. Singh and A. K. Madan, *Application of Graph Theory: Relationship of eccentric Connectivity Index and Wieners Index with Anti-inflammatory Activity*. J. Math. Anal. Appl, **266** (2002), 259–268.
- [13] I. Gutman, N. Trinajstić, *Graph theory and molecular orbitals., Total  $\pi$ -electron energy of alternating hydrocarbons*. Chemical Physics Letters, **17** (1972),, 535-538.
- [14] I. Gutman and O. E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer-Verlag, New York (1986).
- [15] Y. Huo, J. B. Lio, A. Q. Baig, W. Sajjad and M. H. Farahani, *Connective eccentric Index of  $NA_m^n$  Nanotube*. Journal of Computational and Theoretical Nanoscience, **14** (2017), 1832 – –1836.
- [16] H. Hosoya, *Topological index. Newly proposed quantity characterizing the topological nature of structure of isomers of saturated hydrocarbons*. Bull. Chem. Soc. Jpn. , **44** (1971), 2332–2337.
- [17] H. Hosoya, *Topological index as strong sorting device for coding chemical structure*. J. Chem. Doc., **12** (1972), 181–183.
- [18] A. Ilic , I. Gutman, *Eccentric connectivity index of chemical trees*. MATCH Commun Math Comput Chem., , **65(3)** (2011), 731–744.
- [19] M. Randić, *On characterization of molecular branching*. J. Am. Chem. Soc., **97** (1975), 6609–6615.
- [20] V. Sharma, R. Goswami, A.K. Madan, *Eccentric connectivity index: A novel highly discriminating topological descriptor for structure-property and structure-activity studies*. J. Chem. Inf. Comput. Sci. , **37** (1997), 273–282.
- [21] M. K. Siddiqui, M. Naeem, N. A. Rahman, M. Imran, *Computing topological indices of certain networks*. J. of Optoelectronics and Advanced Material, **Vol. 18, No. 9-10** (2016), p. 884 – 892.
- [22] M. Stefu, M. V. Diudea, *Wiener Index of  $C_4C_8$  Nanotubes*. MATCH Commun Math Comput Chem, **50** (2004), 133–144.

- [23] M. H. Sunilkumar, B. K. Bhagyashri, G. B. Ratnamma, and M. G. Vijay, QSPR analysis of certain graph theoretical matrices and their corresponding energy, *Applied Mathematics and Nonlinear Sciences*, vol. 2, no. 1, pp. 131-150, 2017.
- [24] H. Wiener, *Correlation of Heats of Isomerization, and Differences in Heats of Vaporization of Isomers, Among the Paraffin Hydrocarbons* *J. Am. Chem. Soc.* , **69** (1947), 2636–2638.
- [25] H. Wiener, *Influence of interatomic forces on paraffin properties.* *J. Chem. Phys.* , **15** (1947), 766–767.
- [26] H. Wiener, *Vapour-pressure-temperature relations among the branched paraffin hydrocarbons.* *J. Chem. Phys.* , **15** (1948), 425–430.
- [27] B. Zhou, Z. Du , *On eccentric connectivity index.* *MATCH Commun Math Comput Chem* , **63** (2010), 181–198.

<sup>1</sup> DEPARTMENT OF MATHEMATICS, COMSATS INSTITUTE OF INFORMATION TECHNOLOGY, SAHIWAL, PAKISTAN.

*E-mail address:* naeempkn@gmail.com, kamransiddiqui75@gmail.com

<sup>2</sup> DEPARTAMENTO DE MATEMÁTICA APLICADA Y ESTADÍSTICA. UNIVERSIDAD POLITÉCNICA DE CARTAGENA, HOSPITAL DE MARINA, 30203-CARTAGENA, REGIÓN DE MURCIA, SPAIN.–CORRESPONDING AUTHOR–

*E-mail address:* juan.garcia@upct.es

<sup>3</sup>SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, YUNNAN NORMAL UNIVERSITY, KUNMING, CHINA

*E-mail address:* gaowei@ynnu.edu.cn