

Juan L. G. Guirao*, Muhammad Kamran Siddiqui, and Asif Hussain

On Topological Properties of Block Shift and Hierarchical Hypercube Networks

DOI: DOI

Received October, 25 2018; revised ..; accepted ..

Abstract: Networks play an important role in electrical and electronic engineering. It depends on what area of electrical and electronic engineering, for example there is a lot more abstract mathematics in communication theory and signal processing and networking etc. Networks involve nodes communicating with each other. Graph theory has found a considerable use in this area of research. A topological index is a real number associated with chemical constitution purporting for correlation of chemical networks with various physical properties, chemical reactivity. The concept of hyper Zagreb index, first multiple Zagreb index, second multiple Zagreb index and Zagreb polynomials were established in chemical graph theory based on vertex degrees. In this paper, we extend this study to interconnection networks and derive analytical closed results of hyper Zagreb index, first multiple Zagreb index, second multiple Zagreb index, Zagreb polynomials and redefined Zagreb indices for block shift network ($BSN - 1$) and ($BSN - 2$), Hierarchical hypercube ($HHC - 1$) and ($HHC - 2$).

Keywords: hyper Zagreb index, first multiplicative Zagreb index, second multiplicative Zagreb index, Zagreb polynomials, redefined Zagreb indices, block shift networks, Hierarchical interconnection networks.

PACS: 02.10.Ox

*Corresponding Author: Juan L. G. Guirao: Departamento de Matemática Aplicada y Estadística, Universidad Politécnica de Cartagena, Hospital de Marina, 30203-Cartagena, Región de Murcia, Spain, E-mail: juan.garcia@upct.es

Muhammad Kamran Siddiqui: Department of Mathematics, COMSATS University Islamabad, Sahiwal Campus, 57000, Pakistan, E-mail: kamransiddiqui75@gmail.com

Asif Hussain: School of Management and Economics, North China University of Water Resource and Electric Power, Zhengzhou 450046, Henan, P.R. China, E-mail: asifh070@gmail.com

1 Introduction

Multiprocessor interconnection networks are often required to connect thousands of homogeneously replicated processor-memory pairs, each of which is called a processing node. Instead of using a shared memory, all synchronization and communication between processing nodes for program execution is often done via message passing. Design and use of multi processor interconnection networks have recently drawn considerable attention due to the availability of inexpensive, powerful microprocessors and memory chips. The mesh networks have been recognized as versatile interconnection networks for massively parallel computing. Mesh/torus-like low-dimensional networks have recently received a lot of attention for their better scalability to larger networks, as opposed to more complex networks such as hypercubes. In particular the failure of cooperation on dependent networks has been studied a lot recently in [1].

A number of Hierarchical Inter connection network (HIN) provide a framework for designing networks with reduced link cost by designing networks with reduced link cost by taking advantage of the locality of communication that exist in parallel applications. HIN employ multiple level. Lower level network provide local communication while higher level networks facilitate remote communication. The multistage networks have long been used as communication networks for parallel computing [2].

The topological properties of certain networks are studied in [3]. Molecules and molecular compounds are often modeled by molecular graphs. A molecular graph is a graph in which vertices are atoms of a given molecule and edges are its chemical bonds. Since the valency of carbon is four, it is natural to consider all graphs with maximum degree ≤ 4 , as a molecular graph. A graph $G(V, E)$ with vertex set V and edge set E is connected, if there exists a connection between any pair of vertices in G . A network is simply a connected graph having no multiple edges and no loops. Throughout in this article, the degree of a vertex $v \in V(G)$, denoted by $deg(v)$, is the number of edges incident to v .

A *topological index* is a numeric quantity associated with a graph which characterizes the topology of graph and is invariant under graph automorphism. In more precise way, a topological index $Top(G)$ of a graph G , is a number with the property that for every graph H isomorphic to graph G , $Top(H) = Top(G)$. The concept of topological index came from work done by Wiener [4] while he was working on boiling point of paraffin. He named this index as *path number*. Later on, the path number was renamed as *Wiener index*. The Wiener index is the first and the most studied topological index, both from theoretical point of view and applications, and defined as the sum of distances between all pairs of vertices in G , see for details [5, 6].

One of the oldest topological index is the first Zagreb index introduced by I. Gutman and N. Trinajstić based on degree of vertices of G in 1972 [7]. The first and second Zagreb indices of a graph G are defined as:

$$M_1(G) = \sum_{uv \in E(G)} [deg(u) + deg(v)] \quad (1)$$

$$M_2(G) = \sum_{uv \in E(G)} [deg(u) \times deg(v)] \quad (2)$$

In 2013, G. H. Shirdel, H. R. Pour and A. M. Sayadi [8] introduced a new degree based of Zagreb index named "hyper Zagreb index" as:

$$HM(G) = \sum_{uv \in E(G)} [deg(u) + deg(v)]^2 \quad (3)$$

M. Ghorbani and N. Azimi defined two new versions of Zagreb indices of a graph G in 2012 [9]. These indices are the first multiplicative Zagreb index $PM_1(G)$ and the second multiplicative Zagreb index $PM_2(G)$ and these indices are defined as:

$$PM_1(G) = \prod_{uv \in E(G)} [deg(u) + deg(v)] \quad (4)$$

$$PM_2(G) = \prod_{uv \in E(G)} [deg(u) \times deg(v)] \quad (5)$$

The properties of $PM_1(G)$ and $PM_2(G)$ indices for some chemical structures have been studied in [10].

The first Zagreb Polynomial $M_1(G, x)$ and the second Zagreb polynomial $M_2(G, x)$ are defined as:

$$M_1(G, x) = \sum_{uv \in E(G)} x^{deg(u)+deg(v)} \quad (6)$$

$$M_2(G, x) = \sum_{uv \in E(G)} x^{deg(u) \times deg(v)} \quad (7)$$

The properties of Zagreb polynomials for some chemical structures have been studied in [11].

Ranjini et al. [12] reclassified the Zagreb indices to be specific the redefined first, second and third Zagreb indices for a graph G as;

$$ReZG_1(G) = \sum_{uv \in E(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \quad (8)$$

$$ReZG_2(G) = \sum_{uv \in E(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \quad (9)$$

$$ReZG_3(G) = \sum_{uv \in E(G)} \left[[deg(u) \times deg(v)] \times [deg(u) + deg(v)] \right] \quad (10)$$

Nowadays there is an extensive research activity on $HM(G)$, $PM_1(G)$, $PM_2(G)$ indices and $M_1(G, x)$, $M_2(G, x)$ polynomials and their variants, see also [13–15].

For further study of topological indices of various graph families, see [16–30].

2 Methods


For the calculation of our outcomes, we used a methodology for combinatorial enlisting, a vertex segment procedure, an edge parcel procedure, diagram theoretical instruments, logical frameworks, a degree-tallying technique, and a degrees of neighbors system.

3 Results for Block Shift Network


$$(BSN - 1)_{n \times n}$$

In this section, we compute certain degree based topological indices for block shift network (BSN) see Figure 1(a). The Randić type indices for Hierarchical interconnection networks is computed by Haider et. al in [31]. We compute first and second Zagreb indices, hyper Zagreb index $HM(G)$, first multiple Zagreb index $PM_1(G)$, second multiple Zagreb index $PM_2(G)$ and Zagreb polynomials $M_1(G, x)$, $M_2(G, x)$ for $(BSN - 1)_{n \times n}$.

Let G be a block shift network. The number of vertices and edges in $(BSN - 1)_{n \times n}$ are $16n^2$ and $24n^2 - 2$ respectively. There are two types of edges in $(BSN - 1)_{n \times n}$ based on degrees of end vertices of each edge. The edge set of $(BSN - 1)_{n \times n}$ can be divided into two partitions based on the degree of end vertices. The first edge



(a)



(b)

Fig. 1. (a) block shift network $(BSN - 1)_{3 \times 3}$
 (b) block shift network $(BSN - 2)_{1 \times 1}$.

partition $E_1((BSN - 1)_{n \times n})$ contains 8 edges uv , where $deg(u) = 2, deg(v) = 3$. The second edge partition $E_2((BSN - 1)_{n \times n})$ contains $24n^2 - 10$ edges uv , where $deg(u) = deg(v) = 3$.

– **The first and second Zagreb indices of $(BSN - 1)_{n \times n}$**

Now using equations (1),(2), we have

$$\begin{aligned} M_1(G) &= \sum_{uv \in E(G)} [deg(u) + deg(v)] \\ M_1(G) &= \sum_{uv \in E_1(G)} [deg(u) + deg(v)] \\ &+ \sum_{uv \in E_2(G)} [deg(u) + deg(v)] \\ &= 5|E_1(G)| + 6|E_2(G)| \\ &= 5(8) + 6(24n^2 - 10) \\ &= 144n^2 - 20. \end{aligned}$$

$$\begin{aligned} M_2(G) &= \sum_{uv \in E(G)} [deg(u) \times deg(v)] \\ M_2(G) &= \sum_{uv \in E_1(G)} [deg(u) \times deg(v)] \\ &+ \sum_{uv \in E_2(G)} [deg(u) \times deg(v)] \\ &= 6|E_1(G)| + 9|E_2(G)| \\ &= 6(8) + 9(24n^2 - 10) \\ &= 216n^2 - 42. \end{aligned}$$

– **Hyper Zagreb index of $(BSN - 1)_{n \times n}$**

The Hyper Zagreb index using equation (3) is computed as:

$$\begin{aligned} HM(G) &= \sum_{uv \in E(G)} [deg(u) + deg(v)]^2 \\ HM(G) &= \sum_{uv \in E_1(G)} [deg(u) + deg(v)]^2 \\ &+ \sum_{uv \in E_2(G)} [deg(u) + deg(v)]^2 \\ &= 5^2|E_1(G)| + 6^2|E_2(G)| \\ &= 25(8) + 36(24n^2 - 10) \\ &= 864n^2 - 160. \end{aligned}$$

– **Multiplicative Zagreb indices of $(BSN - 1)_{n \times n}$**

The multiplicative Zagreb indices using equations (4), (5) are computed as:

$$\begin{aligned} PM_1(G) &= \prod_{uv \in E(G)} [deg(u) + deg(v)] \\ PM_1(G) &= \prod_{uv \in E_1(G)} [deg(u) + deg(v)] \\ &\times \prod_{uv \in E_2(G)} [deg(u) + deg(v)] \\ &= 5^{|E_1(G)|} \times 6^{|E_2(G)|} \\ &= 5^8 \times 6^{(24n^2 - 10)}. \end{aligned}$$

$$\begin{aligned} PM_2(G) &= \prod_{uv \in E(G)} [deg(u) \times deg(v)] \\ PM_2(G) &= \prod_{uv \in E_1(G)} [deg(u) \times deg(v)] \\ &\times \prod_{uv \in E_2(G)} [deg(u) \times deg(v)] \\ &= 6^{|E_1(G)|} \times 9^{|E_2(G)|} \\ &= 6^8 \times 9^{(24n^2 - 10)}. \end{aligned}$$

– **The first and second Zagreb polynomials of $(BSN - 1)_{n \times n}$**

Now using equations (6),(7), we have

$$\begin{aligned} M_1(G, x) &= \sum_{uv \in E(G)} x^{[deg(u)+deg(v)]} \\ M_1(G, x) &= \sum_{uv \in E_1(G)} x^{[deg(u)+deg(v)]} \\ &+ \sum_{uv \in E_2(G)} x^{[deg(u)+deg(v)]} \\ &= \sum_{uv \in E_1(G)} x^5 + \sum_{uv \in E_2(G)} x^6 \\ &= |E_1(G)|x^5 + |E_2(G)|x^6 \\ &= 8x^5 + (24n^2 - 10)x^6. \end{aligned}$$

$$\begin{aligned} M_2(G, x) &= \sum_{uv \in E(G)} x^{[deg(u) \times deg(v)]} \\ M_2(G, x) &= \sum_{uv \in E_1(G)} x^{[deg(u) \times deg(v)]} \\ &+ \sum_{uv \in E_2(G)} x^{[deg(u) \times deg(v)]} \\ &= \sum_{uv \in E_1(G)} x^6 + \sum_{uv \in E_2(G)} x^9 \\ &= |E_1(G)|x^6 + |E_2(G)|x^9 \\ &= 8x^6 + (24n^2 - 10)x^9. \end{aligned}$$

– **The redefine first, second, and third Zagreb index of $(BSN - 1)_{n \times n}$**

Now using the edge partition of the block shift network G , and equation (8), we have:

$$\begin{aligned} ReZG_1(G) &= \sum_{uv \in E(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \\ &= \sum_{uv \in E_1(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \\ &+ \sum_{uv \in E_2(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \\ &= 8 \left(\frac{2+3}{2 \times 3} \right) + (24n^2 - 10) \left(\frac{3+3}{3 \times 3} \right) \\ &= \frac{48n^2}{3}. \end{aligned}$$

By using equation (9), the second redefine Zagreb index is computed as below:

$$\begin{aligned} ReZG_2(G) &= \sum_{uv \in E(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \\ &= \sum_{ab \in E_1(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \\ &+ \sum_{ab \in E_2(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \\ &= 8 \left(\frac{2 \times 3}{2+3} \right) + (24n^2 - 10) \left(\frac{3 \times 3}{3+3} \right) \\ &= \frac{48}{5} + 36n^2 - 15. \end{aligned}$$

Now by using equation (10), the third redefine Zagreb index is computed as:

$$\begin{aligned}
 ReZG_3(G) &= \sum_{uv \in E(G)} \left[[deg(u) \times deg(v)] \right. \\
 &\quad \times \left. [deg(u) + deg(v)] \right] \\
 &= \sum_{uv \in E_1(G)} \left[[deg(u) \times deg(v)] \right. \\
 &\quad \times \left. [deg(u) + deg(v)] \right] \\
 &+ \sum_{uv \in E_2(G)} \left[[deg(u) \times deg(v)] \right. \\
 &\quad \times \left. [deg(u) + deg(v)] \right] \\
 &= 8 \left((2 \times 3) \times (2 + 3) \right) \\
 &+ (24n^2 - 10) \left((3 \times 3) \times (3 + 3) \right) \\
 &= 1296n^2 - 300.
 \end{aligned}$$

4 Results for Block Shift Network $(BSN - 2)_{n \times n}$

The number of vertices and edges in $(BSN - 2)_{n \times n}$ are $16n^2$ and $32n^2 - 2$ respectively see Figure 1(b). There are two types of edges in $(BSN - 2)_{n \times n}$ based on degrees of end vertices of each edge. The edge set of $(BSN - 2)_{n \times n}$ can be divided into two partitions based on the degree of end vertices. The first edge partition $E_1((BSN - 2)_{n \times n})$ contains 12 edges uv , where $deg(u) = 3, deg(v) = 4$. The second edge partition $E_2((BSN - 2)_{n \times n})$ contains $32n^2 - 14$ edges uv , where $deg(u) = deg(v) = 4$.

– **The first and second Zagreb indices of $(BSN - 2)_{n \times n}$**

Now using equations (1),(2), we have

$$\begin{aligned}
 M_1(G) &= \sum_{uv \in E(G)} [deg(u) + deg(v)] \\
 M_1(G) &= \sum_{uv \in E_1(G)} [deg(u) + deg(v)] \\
 &+ \sum_{uv \in E_2(G)} [deg(u) + deg(v)] \\
 &= 7|E_1(G)| + 8|E_2(G)| \\
 &= 7(12) + 8(32n^2 - 14) \\
 &= 512n^2 + 12.
 \end{aligned}$$

$$\begin{aligned}
 M_2(G) &= \sum_{uv \in E(G)} [deg(u) \times deg(v)] \\
 M_2(G) &= \sum_{uv \in E_1(G)} [deg(u) \times deg(v)] \\
 &+ \sum_{uv \in E_2(G)} [deg(u) \times deg(v)] \\
 &= 12|E_1(G)| + 16|E_2(G)| \\
 &= 12(12) + 16(32n^2 - 14) \\
 &= 512n^2 - 80.
 \end{aligned}$$

– **Hyper Zagreb index of $(BSN - 2)_{n \times n}$**

The Hyper Zagreb index using equation (3) is computed as:

$$\begin{aligned}
 HM(G) &= \sum_{uv \in E(G)} [deg(u) + deg(v)]^2 \\
 HM(G) &= \sum_{uv \in E_1(G)} [deg(u) + deg(v)]^2 \\
 &+ \sum_{uv \in E_2(G)} [deg(u) + deg(v)]^2 \\
 &= 7^2|E_1(G)| + 8^2|E_2(G)| \\
 &= 49(12) + 64(32n^2 - 14) \\
 &= 2048n^2 - 308.
 \end{aligned}$$

– **Multiplicative Zagreb indices of $(BSN - 2)_{n \times n}$**

The multiplicative Zagreb indices using equations (4), (5) are computed as:

$$\begin{aligned} PM_1(G) &= \prod_{uv \in E(G)} [deg(u) + deg(v)] \\ PM_1(G) &= \prod_{uv \in E_1(G)} [deg(u) + deg(v)] \\ &\times \prod_{uv \in E_2(G)} [deg(u) + deg(v)] \\ &= 7^{|E_1(G)|} \times 8^{|E_2(G)|} \\ &= 7^{12} \times 8^{(32n^2 - 14)}. \end{aligned}$$

$$\begin{aligned} PM_2(G) &= \prod_{uv \in E(G)} [deg(u) \times deg(v)] \\ PM_2(G) &= \prod_{uv \in E_1(G)} [deg(u) \times deg(v)] \\ &\times \prod_{uv \in E_2(G)} [deg(u) \times deg(v)] \\ &= 12^{|E_1(G)|} \times 16^{|E_2(G)|} \\ &= 12^{12} \times 16^{(32n^2 - 14)}. \end{aligned}$$

– **The first and second Zagreb polynomials of $(BSN - 2)_{n \times n}$**

Now using equations (6),(7), we have

$$\begin{aligned} M_1(G, x) &= \sum_{uv \in E(G)} x^{[deg(u)+deg(v)]} \\ M_1(G, x) &= \sum_{uv \in E_1(G)} x^{[deg(u)+deg(v)]} \\ &+ \sum_{uv \in E_2(G)} x^{[deg(u)+deg(v)]} \\ &= \sum_{uv \in E_1(G)} x^7 + \sum_{uv \in E_2(G)} x^8 \\ &= |E_1(G)|x^7 + |E_2(G)|x^8 \\ &= 12x^7 + (32n^2 - 14)x^8. \end{aligned}$$

$$\begin{aligned} M_2(G, x) &= \sum_{uv \in E(G)} x^{[deg(u) \times deg(v)]} \\ M_2(G, x) &= \sum_{uv \in E_1(G)} x^{[deg(u) \times deg(v)]} \\ &+ \sum_{uv \in E_2(G)} x^{[deg(u) \times deg(v)]} \\ &= \sum_{uv \in E_1(G)} x^{12} + \sum_{uv \in E_2(G)} x^{16} \\ &= |E_1(G)|x^{12} + |E_2(G)|x^{16} \\ &= 12x^{12} + (32n^2 - 14)x^{16}. \end{aligned}$$

– **The redefine first, second, and third Zegreb index of $(BSN - 2)_{n \times n}$**

Now using the using equation (8) and the edge partition of the block shift network $(BSN - 2)_{n \times n}$, we have:

$$\begin{aligned} ReZG_1(G) &= \sum_{uv \in E(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \\ &= \sum_{ab \in E_1(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \\ &+ \sum_{ab \in E_2(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \\ &= 12 \left(\frac{3+4}{3 \times 4} \right) + (32n^2 - 14) \left(\frac{4+4}{4 \times 4} \right) \\ &= 16n^2. \end{aligned}$$

By using equation (9), the second redefine Zagreb index is computed as below:

$$\begin{aligned} ReZG_2(G) &= \sum_{uv \in E(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \\ &= \sum_{uv \in E_1(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \\ &+ \sum_{uv \in E_2(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \\ &= 12 \left(\frac{3 \times 4}{3+4} \right) + (32n^2 - 14) \left(\frac{4 \times 4}{4+4} \right) \\ &= \frac{144}{7} + 64n^2 - 58. \end{aligned}$$

Now by using equation (10), the third redefine Zagreb index is computed as:

$$\begin{aligned}
 ReZG_3(G) &= \sum_{uv \in E(G)} [deg(u) \times deg(v)] \\
 &\times [deg(u) + deg(v)] \\
 &= \sum_{uv \in E_1(G)} [deg(u) \times deg(v)] \\
 &\times [deg(u) + deg(v)] \\
 &+ \sum_{uv \in E_2(G)} [deg(u) \times deg(v)] \\
 &\times [deg(u) + deg(v)] \\
 &= 12((3 \times 4) \times (3 + 4)) \\
 &+ (32n^2 - 14)((4 \times 4) \times (4 + 4)) \\
 &= 4096n^2 - 2800.
 \end{aligned}$$

5 Results for Hierarchical Hypercube Network

$(HHC - 1)_{n \times n}$

In this section, we compute certain degree based topological indices of Hierarchical interconnection networks see Figure 2(a). The Randic type indices for Hierarchical interconnection networks is computed by Haider et. al in [31]. We compute first and second Zagreb indices, hyper Zagreb index $HM(G)$, first multiple Zagreb index $PM_1(G)$, second multiple Zagreb index $PM_2(G)$ and Zagreb polynomials $M_1(G, x)$, $M_2(G, x)$ for Hierarchical Hypercube network $(HHC - 1)_{n \times n}$ and Hierarchical Hypercube network $(HHC - 2)_{n \times n}$.

The number of vertices and edges in $(HHC - 1)_{n \times n}$ are $16n + 16$ and $24n + 20$ respectively. There are two types of edges in $(HHC - 1)_{n \times n}$ based on degrees of end vertices of each edge. The edge set of $(HHC - 1)_{n \times n}$ can be divided into two partitions based on the degree of end vertices. The first edge partition $E_1((HHC - 1)_{n \times n})$ contains 16 edges uv , where $deg(u) = 2, deg(v) = 3$. The second edge partition $E_2((HHC - 1)_{n \times n})$ contains $24n + 4$ edges uv , where $deg(u) = deg(v) = 3$.

– **The first and second Zagreb indices of $(HHC - 1)_{n \times n}$**




Fig. 2. (a) Hierarchical Hypercube network $(HHC - 1)_{1 \times 1}$
(b) Hierarchical Hypercube network $(HHC - 2)_{1 \times 1}$.

Now using equations (1),(2), we have

$$\begin{aligned}
 M_1((HHC - 1)_{n \times n}) &= \sum_{uv \in E(G)} [deg(u) + deg(v)] \\
 M_1(G) &= \sum_{uv \in E_1(G)} [deg(u) + deg(v)] \\
 &+ \sum_{uv \in E_2(G)} [deg(u) + deg(v)] \\
 &= 5|E_1(G)| + 6|E_2(G)| \\
 &= 5(16) + 6(24n + 4) \\
 &= 216n + 104.
 \end{aligned}$$

$$\begin{aligned}
 M_2(G) &= \sum_{uv \in E(G)} [deg(u) \times deg(v)] \\
 M_2(G) &= \sum_{uv \in E_1(G)} [deg(u) \times deg(v)] \\
 &+ \sum_{uv \in E_2(G)} [deg(u) \times deg(v)] \\
 &= 6|E_1(G)| + 9|E_2(G)| \\
 &= 6(16) + 9(24n + 4) \\
 &= 216n + 132.
 \end{aligned}$$

– **Hyper Zagreb index of $(HHC - 1)_{n \times n}$**

The Hyper Zagreb index using equation (3) is computed as: Now using equations (6),(7), we have

$$\begin{aligned}
 HM(G) &= \sum_{uv \in E(G)} [deg(u) + deg(v)]^2 \\
 HM(G) &= \sum_{uv \in E_1(G)} [deg(u) + deg(v)]^2 \\
 &+ \sum_{uv \in E_2(G)} [deg(u) + deg(v)]^2 \\
 &= 5^2|E_1(G)| + 6^2|E_2(G)| \\
 &= 25(16) + 36(24n + 4) \\
 &= 864n + 544.
 \end{aligned}$$

$$\begin{aligned}
 M_1(G, x) &= \sum_{uv \in E(G)} x^{[deg(u)+deg(v)]} \\
 M_1(G, x) &= \sum_{uv \in E_1(G)} x^{[deg(u)+deg(v)]} \\
 &+ \sum_{uv \in E_2(G)} x^{[deg(u)+deg(v)]} \\
 &= \sum_{uv \in E_1(G)} x^5 + \sum_{uv \in E_2(G)} x^6 \\
 &= |E_1(G)|x^5 + |E_2(G)|x^6 \\
 &= 16x^5 + (24n + 4)x^6.
 \end{aligned}$$

– **Multiple Zagreb indices of $(HHC - 1)_{n \times n}$**

The Multiple-Zagreb indices using equations (4), (5) are computed as:

$$\begin{aligned}
 PM_1(G) &= \prod_{uv \in E(G)} [deg(u) + deg(v)] \\
 PM_1(G) &= \prod_{uv \in E_1(G)} [deg(u) + deg(v)] \\
 &\times \prod_{uv \in E_2(G)} [deg(u) + deg(v)] \\
 &= 5^{|E_1(G)|} \times 6^{|E_2(G)|} \\
 &= 5^{16} \times 6^{(24n+4)}.
 \end{aligned}$$

$$\begin{aligned}
 PM_2(G) &= \prod_{uv \in E(G)} [deg(u) \times deg(v)] \\
 PM_2(G) &= \prod_{uv \in E_1(G)} [deg(u) \times deg(v)] \\
 &\times \prod_{uv \in E_2(G)} [deg(u) \times deg(v)] \\
 &= 6^{|E_1(G)|} \times 9^{|E_2(G)|} \\
 &= 6^{16} \times 9^{(24n+4)}.
 \end{aligned}$$

– **The first and second Zagreb polynomials of $(HHC - 1)_{n \times n}$**

$$\begin{aligned}
 M_2(G, x) &= \sum_{uv \in E(G)} x^{[deg(u) \times deg(v)]} \\
 M_2(G, x) &= \sum_{uv \in E_1(G)} x^{[deg(u) \times deg(v)]} \\
 &+ \sum_{uv \in E_2(G)} x^{[deg(u) \times deg(v)]} \\
 &= \sum_{uv \in E_1(G)} x^6 + \sum_{uv \in E_2(G)} x^9 \\
 &= |E_1(G)|x^6 + |E_2(G)|x^9 \\
 &= 16x^6 + (24n + 4)x^9.
 \end{aligned}$$

– **The redefine first, second, and third Zegreb index of $(HHC - 1)_{n \times n}$**

Now using the using equation (8) and the edge partition of the block shift network $(HHC - 1)_{n \times n}$, we have:

$$\begin{aligned}
 ReZG_1(G) &= \sum_{uv \in E(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \\
 &= \sum_{uv \in E_1(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \\
 &+ \sum_{uv \in E_2(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \\
 &= 16 \left(\frac{3+2}{3 \times 2} \right) + (24n + 4) \left(\frac{3+3}{3 \times 3} \right) \\
 &= \frac{40}{3} + \frac{48n^2 + 8}{3}.
 \end{aligned}$$

By using equation (9), the second redefine Zagreb index is computed as below:

$$\begin{aligned} ReZG_2(G) &= \sum_{uv \in E(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \\ &= \sum_{uv \in E_1(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \\ &+ \sum_{uv \in E_2(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \\ &= 16\left(\frac{3 \times 2}{3 + 2}\right) + (24n + 4)\left(\frac{3 \times 3}{3 + 3}\right) \\ &= \frac{96}{5} + 36n + 6. \end{aligned}$$

Now by using equation (10), the third redefine Zagreb index is computed as:

$$\begin{aligned} ReZG_3(G) &= \sum_{uv \in E(G)} \left[[deg(u) \times deg(v)] \right. \\ &\times \left. [deg(u) + deg(v)] \right] \\ &= \sum_{uv \in E_1(G)} \left[[deg(u) \times deg(v)] \right. \\ &\times \left. [deg(u) + deg(v)] \right] \\ &+ \sum_{uv \in E_2(G)} \left[[deg(u) \times deg(v)] \right. \\ &\times \left. [deg(u) + deg(v)] \right] \\ &= 16\left((3 \times 2) \times (3 + 2)\right) \\ &+ (24n + 4)\left((3 \times 3) \times (3 + 3)\right) \\ &= 1296n + 576. \end{aligned}$$

6 Results for Hierarchical Hypercube Network

$(HHC - 2)_{n \times n}$

The number of vertices and edges in $(HHC - 2)_{n \times n}$ are $16n^2$ and $32n^2 - 2$ respectively see Figure 2(b). There are two types of edges in $(HHC - 2)_{n \times n}$ based on degrees of end vertices of each edge. The edge set of $(HHC - 2)_{n \times n}$ can be divided into two partitions based on the degree of end vertices. The first edge partition $E_1((HHC - 2)_{n \times n})$ contains 24 edges uv , where $deg(u) = 3, deg(v) = 4$. The

second edge partition $E_2((HHC - 2)_{n \times n})$ contains $32n + 4$ edges uv , where $deg(u) = deg(v) = 4$.

– **The first and second Zagreb indices of $(HHC - 2)_{n \times n}$**

Now using equations (1),(2), we have

$$\begin{aligned} M_1(G) &= \sum_{uv \in E(G)} [deg(u) + deg(v)] \\ M_1(G) &= \sum_{uv \in E_1(G)} [deg(u) + deg(v)] \\ &+ \sum_{uv \in E_2(G)} [deg(u) + deg(v)] \\ &= 7|E_1(G)| + 8|E_2(G)| = 7(24) + 8(32n + 4) \\ &= 512n + 200. \end{aligned}$$

$$\begin{aligned} M_2(G) &= \sum_{uv \in E(G)} [deg(u) \times deg(v)] \\ M_2(G) &= \sum_{uv \in E_1(G)} [deg(u) \times deg(v)] \\ &+ \sum_{uv \in E_2(G)} [deg(u) \times deg(v)] \\ &= 12|E_1(G)| + 16|E_2(G)| \\ &= 12(24) + 16(32n + 4) \\ &= 512n + 352. \end{aligned}$$

– **Hyper Zagreb index of $(HHC - 2)_{n \times n}$**

The Hyper Zagreb index using equation (3) is computed as:

$$\begin{aligned} HM(G) &= \sum_{uv \in E(G)} [deg(u) + deg(v)]^2 \\ HM(G) &= \sum_{uv \in E_1(G)} [deg(u) + deg(v)]^2 \\ &+ \sum_{uv \in E_2(G)} [deg(u) + deg(v)]^2 \\ &= 7^2|E_1(G)| + 8^2|E_2(G)| \\ &= 49(24) + 64(32n + 4) \\ &= 2048n + 1432. \end{aligned}$$

– **Multiple Zagreb indices of $(HHC - 2)_{n \times n}$**

The Multiple-Zagreb indices using equations (4), (5) are computed as:

$$\begin{aligned} vPM_1(G) &= \prod_{uv \in E(G)} [deg(u) + deg(v)] \\ PM_1(G) &= \prod_{uv \in E_1(G)} [deg(u) + deg(v)] \\ &\times \prod_{uv \in E_2(G)} [deg(u) + deg(v)] \\ &= 7^{|E_1(G)|} \times 8^{|E_2(G)|} \\ &= 7^{24} \times 8^{(32n+4)}. \end{aligned}$$

$$\begin{aligned} PM_2(G) &= \prod_{uv \in E(G)} [deg(u) \times deg(v)] \\ PM_2(G) &= \prod_{uv \in E_1(G)} [deg(u) \\ &\times deg(v)] \times \prod_{uv \in E_2(G)} [deg(u) \times deg(v)] \\ &= 12^{|E_1(G)|} \times 16^{|E_2(G)|} \\ &= 12^{24} \times 16^{(32n+4)}. \end{aligned}$$

– **The first and second Zagreb polynomials of $(HHC - 2)_{n \times n}$**

Now using equations (6),(7), we have

$$\begin{aligned} M_1(G, x) &= \sum_{uv \in E(G)} x^{[deg(u)+deg(v)]} \\ M_1(G, x) &= \sum_{uv \in E_1(G)} x^{[deg(u)+deg(v)]} \\ &+ \sum_{uv \in E_2(G)} x^{[deg(u)+deg(v)]} \\ &= \sum_{uv \in E_1(G)} x^7 + \sum_{uv \in E_2(G)} x^8 \\ &= |E_1(G)|x^7 + |E_2(G)|x^8 \\ &= 24x^7 + (32n + 4)x^8. \end{aligned}$$

$$\begin{aligned} M_2(G, x) &= \sum_{uv \in E(G)} x^{[deg(u) \times deg(v)]} \\ M_2(G, x) &= \sum_{uv \in E_1(G)} x^{[deg(u) \times deg(v)]} \\ &+ \sum_{uv \in E_2(G)} x^{[deg(u) \times deg(v)]} \\ &= \sum_{uv \in E_1(G)} x^{12} + \sum_{uv \in E_2(G)} x^{16} \\ &= |E_1(G)|x^{12} + |E_2(G)|x^{16} \\ &= 24x^{12} + (32n + 4)x^{16}. \end{aligned}$$

– **The redefine first, second, and third Zagreb index of $(HHC - 2)_{n \times n}$**

Now using the using equation (8) and the edge partition of the block shift network $(HHC - 2)_{n \times n}$, we have:

$$\begin{aligned} ReZG_1(G) &= \sum_{uv \in E(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \\ &= \sum_{uv \in E_1(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \\ &+ \sum_{uv \in E_2(G)} \frac{deg(u) + deg(v)}{deg(u) \times deg(v)} \\ &= 24 \left(\frac{3+4}{3 \times 4} \right) + (32n + 4) \left(\frac{4+4}{4 \times 4} \right) \\ &= 16n + 16. \end{aligned}$$

By using equation (9), the second redefine Zagreb index is computed as below:

$$\begin{aligned} ReZG_2(G) &= \sum_{uv \in E(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \\ &= \sum_{uv \in E_1(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \\ &+ \sum_{uv \in E_2(G)} \frac{deg(u) \times deg(v)}{deg(u) + deg(v)} \\ &= 24 \left(\frac{3 \times 4}{3 + 4} \right) + (32n + 4) \left(\frac{4 \times 4}{4 + 4} \right) \\ &= \frac{288}{7} + 64n + 8. \end{aligned}$$

Now by using equation (10), the third redefine Zagreb index is computed as:

$$\begin{aligned}
 ReZG_3(G) &= \sum_{uv \in E(G)} \left[[deg(u) \times deg(v)] \right. \\
 &\times [deg(u) + deg(v)] \left. \right] \\
 &= \sum_{uv \in E_1(G)} \left[[deg(u) \times deg(v)] \right. \\
 &\times [deg(u) + deg(v)] \left. \right] \\
 &+ \sum_{uv \in E_2(G)} \left[[deg(u) \times deg(v)] \right. \\
 &\times [deg(u) + deg(v)] \left. \right] \\
 &= 24 \left((3 \times 4) \times (3 + 4) \right) \\
 &+ (32n + 4) \left((4 \times 4) \times (4 + 4) \right) \\
 &= 4608n + 2596.
 \end{aligned}$$

Conclusion

In this paper we determined first Zagreb index $M_1(G)$, second Zagreb index $M_2(G)$, hyper Zagreb index $HM(G)$, first multiple Zagreb index $PM_1(G)$, second multiple Zagreb index $PM_2(G)$, Zagreb polynomials $M_1(G, x)$ and $M_2(G, x)$ and redefined Zagreb indices for block shift networks and Hierarchical Hypercube networks. In future, we are interested in designing some new architectures/networks and then studying their topological indices which will be quite helpful to understand their underlying topologies.

Acknowledgement

The authors are grateful to the anonymous referees for their valuable comments and suggestions that improved this paper.

References

- [1] Gutman I., Tošović J., Radenković S., Marković S., On atom–bond connectivity index and its chemical applicability, *Indian J. Chem.*, 2012, 51A, 690-694.
- [2] Konstantinidou, S., The selective extra stage butterfly, *IEEE. Trans., Very Large Scale Integr. (VLSI) Syst.*, 1992, 1, 502–506.
- [3] Shao Z., Siddiqui M. K., Muhammad M. H., Computing Zagreb Indices and Zagreb Polynomials for Symmetrical Nanotubes. *Symmetry*, 2018, 10, 1-20.
- [4] Wiener H., Structural determination of paraffin boiling points, *J. Am. Chem. Soc.*, 1947, 69, 17-20.
- [5] Dobrynin A.A., Entringer R., Gutman I., Wiener index of trees: Theory and applications, *Acta Appl. Math.*, 2001, 66, 211-249.
- [6] Gutman I., Polansky O. E., *Mathematical Concepts in Organic Chemistry*, Springer-Verlag, New York, 1986.
- [7] Gutman I., Trinajstić N., Graph theory and molecular orbitals total π -electron energy of alternant hydrocarbons, *Chem. Phys. Lett.*, 1972, 17, 535–538.
- [8] Shirdel G.H., Pour H.R., Sayadi A.M., The Hyper-Zagreb Index of Graph Operations, *Iran. J. Math. Chem.*, 2013, 4(2), 213-220.
- [9] Ghorbani, M., Azimi, N., Note on multiple Zagreb indices, *Iran. J. Math. Chem.*, 2012, 32, 137-143.
- [10] Eliasi M., Iranmanesh A., Gutman I., Multiplicative versions of first Zagreb Index, *MATCH Commun. Math. Comput. Chem.*, 2012, 68, 217-230.
- [11] Gutman I., Das K.C., Some Properties of the Second Zagreb Index, *MATCH Commun. Math. Comput. Chem.*, 2004, 50, 103-112.
- [12] Ranjini P.S., Lokesh V., Usha A., Relation between phenylene and hexagonal squeeze using harmonic index, *Int. J. Graph. Theory.*, 2013, 1, 116-21.
- [13] Gao W., Siddiqui M.K., Naeem M., Imran M., Computing multiple ABC index and multiple GA index of some grid graphs, *Open Phys.*, 2018, 161, 588-598.
- [14] Liu J. B., Siddiqui M.K., Zahid M. A., Naeem M., Baig A.Q., Topological Properties of Crystallographic Structure of Molecules. *Symmetry*, 2018, 10, 1-21.
- [15] Siddiqui M. K., Gharibi W., On Zagreb Indices, Zagreb Polynomials of Mesh Derived Networks, *J. Comput. Theor. Nanosci.*, 2016, 13, 8683-8688.
- [16] Gao W., Siddiqui M.K., Naeem M., Rehman N.A., Topological Characterization of Carbon Graphite and Crystal Cubic Carbon Structures, *Molecules*, 2017, 22, 1496-1507.
- [17] Imran M., Siddiqui M. K., Naeem M., Iqbal M. A., On Topological Properties of Symmetric Chemical Structures, *Symmetry*, 2018, 10, 1-20.
- [18] Siddiqui M. K., Naeem M., Rahman N. A., Imran M., Computing topological indices of certain networks, *J. Optoelectron. Adv. Mater.*, 2016, 18, 884-892.
- [19] Gao W., Siddiqui M.K., Molecular Descriptors of Nanotube, Oxide, Silicate, and Triangulene Networks, *J. Chem.*, 2017, 2017, 654-664.
- [20] Basavanagoud B., Gao W., Patil S., Desai V.R., Mirajkar K.G., Pooja, B., Computing First Zagreb index and F-index of New C-products of Graphs, *Applied Mathematics and Nonlinear Sciences*, 2017, 21, 285-298.
- [21] Basavanagoud B., Desai V.R., Patil S., (β, α) Connectivity Index of Graphs, *Applied. Math. Nonlin. Sci.*, 2017, 21, 21-30.

- [22] Gao W., Siddiqui M.K., Imran M., Jamil M.K., Farahani M.R., Forgotten Topological Index of Chemical Structure in Drugs, Saudi Pharm. Jou., 2016, 24, 258-267.
- [23] Gharibi W., Ahmad A., Siddiqui M.K., On Zagreb Indices, Zagreb Polynomials of Nanocone and Nanotubes, J. Comput. Theor. Nanosci., 2016, 13, 5086-5092.
- [24] Idrees N., Naeem M.N., Hussain F., Sadiq A., Siddiqui M.K., Molecular Descriptors of Benzenoid System, Quimica Nova, 2017, 40(2), 143-145.
- [25] 2. Kang S.M., Siddiqui M.K., Rehman N.A., Naeem M., Muhammad M.H., Topological Properties of 2-Dimensional Silicon-Carbons" IEEE Access, 2018, 6(1), 59362-59373.
- [26] Gao W., Imran M., Siddiqui M.K., Naeeme M., Jamil F., Molecular Description of Copper (I) Oxide and Copper (II) Oxide, Quimica. Nova. 2018, 41(8), 874-879.
- [27] 3. Imran M., Siddiqui M.K., Ahmad A., Ali U., Hanif N., On the Degree-Based Topological Indices of the Tickysim SpiNNaker Model, Axioms, 2018, 7, 1-13.
- [28] Randić M., Characterization of molecular branching, J. Am. Chem. Soc., 1975, 97 6609-6615.
- [29] Ramane H.S., Jummannaver R.B., Note on forgotten topological index of chemical structure in drugs, Appl. Math. Nonlinear Sciences., 2016, 12, 369-374.
- [30] Siddiqui M.K., Imran M., Ali A., On Zagreb indices, Zagreb polynomials of some nanostar dendrimers, Appl. Math. Comput., 2016, 280, 132-139.
- [31] Ali H., Baig A.Q., Shafiq M.K., On topological properties of hierarchical interconnection networks, J. Appl. Math. Comput., 2017, 55, 313-334.